

ESO Phase 3 Data Release Description

| | |
|------------------------|-------------|
| Data Collection | VIDEO_XMM |
| Release Number | 2 |
| Data Provider | Matt Jarvis |
| Date | 17.12.2013 |

Abstract

VIDEO is a deep near-infrared survey which targets ~ 12 square degrees over the ELAIS-S1, XMMLSS, and ECDFS extragalactic fields. This document describes the release of the deep stacked data in the VIDEO-XMM3 field in Z,Y,J,H and Ks bands. To the 5σ limit, our Ks catalogue contains 198178 sources over ~ 1.8 deg².

Overview of Observations

We release the individual tiles and associated sources lists along with deep stacked images and associated catalogues separately for each band, along with a Ks-band selected catalogue from the VISTA Deep Extragalactic Observations (VIDEO) Survey. The data includes Z,Y,J,H,Ks deep-stacks and catalogues in a single VISTA tile (~ 1.8 degrees²) over the VIDEO-XMM3 tile. We also release individual tiles in Z and Y bands in the XMM1 field and Y,J and Ks bands in the XMM2 field, along with the associated source lists (see Fig. 1) taken between 01/10/2010 - 31/12/2011. The individual tiles, source lists and deep-stack data over the VIDEO-XMM3 field are fully described in the VIDEO Survey paper (Jarvis et al., 2013) with observations acquired between 03/11/2009 and 20/11/2011.

Release Content

We release individual tile data in Z,Y,J,H and Ks bands over the full XMMLSS survey region (Figure 1), along with a single deep-stacked VISTA tile centred on 2h 26m 18s -04s 44m 00s (VIDEO-XMM3 in Z,Y,J,H,Ks filters).

We also provide a Ks-selected catalogue "VIDEO_XMM3_Kselected_merged_DR1.fits" that provides band-merged photometry using the Ks-band as the detection image and extracting photometry from the shorter wavelength imaging data. We recommend using only those sources with $K_s < 23.5$ for most studies as this gives a completeness of ~ 90 per cent (Jarvis et al. 2013). In the Ks-selected band-merged catalogue, columns where the value is "NaN" denote that the object was detected at the position of the Ks-band source with negative flux, this occurs over all apertures and filters for different objects, but is mostly a problem for the larger aperture.

In addition to the XMM3 YJHKs tiles and associated source lists which were released in DR1 we include further tiles in XMM3, along with the new complete Z-band tile set. We also release single-OB tiles in ZY over XMM1 and YJKs single-OB tiles over XMM2.




Figure. 1 The coordinates of the VIDEO-XMM field. This release comprises data over all three tiles. The deep stack data is only released over the XMM3 tile.

Table 1. Individual tiles released, including data releases DR1 and DR2. Note that two observing strategies for the Z-band data in the VIDEO-XMM3 field were used. These are denoted as Z(1) and Z(2). These can be differentiated between within the image headers by checking the EXPTIME keywords, which corresponds to the *Integration/pixel* in the table below. We note that in a few cases Z(1) is not 2970 due to the non-completion of OBs. The number in brackets in column 5 denotes the number of tiles in the total that were released as part of DR1.

| Field | Filter | <i>Integration/tile</i> | <i>Integration/pixel</i> | <i>Number of Tiles</i> |
|------------|--------|-------------------------|--------------------------|------------------------|
| VIDEO-XMM1 | Z | 2700 sec | 900 sec | 7 |
| VIDEO-XMM1 | Y | 2880 sec | 960 sec | 32 |
| VIDEO-XMM2 | Y | 2880 sec | 960 sec | 12 |
| VIDEO-XMM2 | J | 2880 sec | 960 sec | 8 |
| VIDEO-XMM2 | Ks | 2520 sec | 840 sec | 11 |
| VIDEO-XMM3 | Z(1) | 2970 sec | 990 sec | 17 |
| VIDEO-XMM3 | Z(2) | 3000 sec | 1000 sec | 50 |
| VIDEO-XMM3 | Y | 2880 sec | 960 sec | 28 (23) |
| VIDEO-XMM3 | J | 2880 sec | 960 sec | 34 (29) |
| VIDEO-XMM3 | H | 2520 sec | 840 sec | 26 (21) |
| VIDEO-XMM3 | Ks | 2520 sec | 840 sec | 28 (24) |

Table 2. Measured 5σ depths for the VIDEO-XMM deep-stack release in 1-5 arcsec diameter apertures in AB magnitudes. The time given is the total integration time of the paw-prints used to measure these values.

| Filter | Time (h) Per pixel | 1" (5σ) | 2" (5σ) | 3" (5σ) | 4" (5σ) | 5" (5σ) |
|--------|-----------------------|---------------------|---------------------|---------------------|---------------------|---------------------|
| Z | 17.13 | 26.89 | 25.66 | 24.87 | 24.33 | 23.83 |
| Y | 6.05 | 25.60 | 24.51 | 23.74 | 23.18 | 22.73 |
| J | 9.65 | 25.59 | 24.44 | 23.70 | 23.11 | 22.63 |
| H | 8.13 | 25.25 | 24.12 | 23.42 | 22.87 | 22.40 |
| Ks | 9.02 | 24.86 | 23.77 | 23.10 | 22.53 | 22.13 |

Release Notes

Data Reduction and Calibration

Initial data reduction steps are performed at the Cambridge Astronomical Survey Unit (CASU) using a software pipeline developed specifically for the reduction of VIRCAM data, as part of the VISTA Data Flow System described by Irwin et al. (2004) and updated on the CASU webpages. This pipeline is modular, and allows different processing recipes to be applied to data obtained with different observing strategies.

For the VIDEO survey, the following steps are applied to each raw data frame (itself the result of NDIT double-correlated- sampling exposures which are co-added by the data acquisition system):

- Destriping – removes a low-level horizontal stripe pattern introduced by the controller and correlates over the four detectors on each controller,
- linearity correction – corrects for the non-linear detector response which is typically 2-4 per cent depending on the detector,
- dark correction – subtracts a mean dark exposure from the image, correcting for dark current and some other electronic effects,
- flat field correction – divides images by a mean twilight sky image, to correct for position-dependent variations in telescope/camera throughput and detector response,
- sky background correction – subtracts an estimate of the atmospheric emission (more details can be found in Jarvis et al. 2013, MNRAS, 428, 1281),
- astrometric calibration – compares the positions of stars in the image with those in the 2MASS point source catalogue (Skrutskie et al. 2006),
- photometric calibration – calculates magnitudes in the VISTA Vega-magnitude photometric system for unsaturated 2MASS stars in the image using their magnitudes and colours in the 2MASS point source catalogue, to set the zero point in the VISTA photometric system, and
- jitter stacking – combines the jittered images in a single paw- print position, using a mean stack of bi-linearly-resampled images, with outlier rejection to remove cosmic rays, fast-moving objects, and bad pixels.

The stacked images provided are based on the paw-print data and use only those paw-print images with <0.9 arcsec seeing. As such parts of completed OBs are not included in the deep stacks, on the other hand some data from aborted OBs are still incorporated in the deep stacks if the individual paw-print met the seeing constraint. Due to the independent data selection for the deep images and the single-OB images, it results for example that the deep co-added Ks image of the XMM3 field features a total integration time per pixel that is higher by ~ 2.5 hours than the arithmetic sum of the integration times of the individual Ks tiles (~ 6.5 h). In this case data from in total 44 OBs has been included in the deep image (though not entirely), while the number of single-OB tiles amounts to 28.

For the source catalogue:

- Source detection was performed separately for each individual filter. For the deep-stack single-band source lists and the band-merged source catalogue, we carry out the detection and measurement of source photometry using SExtractor (Bertin & Arnouts 1996),

with the following parameters (see SExtractor documentation for more information):

- DETECT_MINAREA 3
- THRESH_TYPE RELATIVE
- DETECT_THRESH 3
- ANALYSIS_THRESH 3
- FILTER Y
- FILTER_NAME default.conv
- DEBLEND_NTHRESH 32
- DEBLEND_MINCONT 0.0008
- CLEAN Y
- CLEAN_PARAM 1.0
- MASK_TYPE CORRECT
- PHOT_AUTOPARAMS 2.5, 3.5
- PHOT_PETROPARAMS 2.0, 3.5
- BACK_TYPE AUTO
- BACK_SIZE 64
- BACK_FILTERSIZE 3
- BACKPHOTO_TYPE LOCAL
- BACKPHOTO_THICK 24
- BACK_FILTTHRESH 0.0
- Astrometry is referenced to 2MASS.
- Photometry is performed in the Vega system, and is calibrated by reference to 2MASS stars, using the following colour equations (taken from <http://casu.ast.cam.ac.uk/surveys-projects/vista/technical/photometric-properties>):
 - $Z_{\text{VIRCAM}} = J_{2\text{MASS}} + 1.025*(J-H)_{2\text{MASS}}$
 - $Y_{\text{VIRCAM}} = J_{2\text{MASS}} + 0.610*(J-H)_{2\text{MASS}}$
 - $J_{\text{VIRCAM}} = J_{2\text{MASS}} - 0.077*(J-H)_{2\text{MASS}}$
 - $H_{\text{VIRCAM}} = H_{2\text{MASS}} + 0.032*(J-H)_{2\text{MASS}}$
 - $Ks_{\text{VIRCAM}} = Ks_{2\text{MASS}} + 0.010*(J-Ks)_{2\text{MASS}}$
- The released catalogue are on the AB magnitude system with the following used to convert from Vega to AB.
 $Z(+0.521)$, $Y(+0.618)$, $J(+0.937)$, $H(+1.384)$, $Ks(+1.839)$
- No illumination correction has been applied.
- No correction for extinction has been applied. However, extinction is extremely low for these high-galactic-latitude fields.

For the individual tiles the associated tile-specific source lists are created using the CASU VSA pipeline, details of which can be found at <http://casu.ast.cam.ac.uk/surveys-projects/vista/technical/tiles>

Data Quality

- Astrometric checks have been carried out by comparing with the 2MASS stars within the tiled fields. No systematic offsets are found and the rms scatter is found to be ~ 0.25 arcsec.
- The zero point is uniform across the tile to within the quoted uncertainty (< 0.01 mag).
- Photometry has also been checked against unresolved 2MASS sources within the VIDEO field. We find very good agreement between 2MASS and VIDEO after applying the colour equations to move from the 2MASS system to the VIDEO photometric system.
- The fraction of spurious sources, calculated by inverting the tiled image and re-extracting the sources using the same prescription used for the real extraction, is 0.07 per cent for the Ks-band selected catalogue to the 5sigma depth of 23.77 (2arcsec aperture).
- Table 3 in Jarvis et al. (2013) provides the completeness for the various filters over the VIDEO-XMM3 deep-stack.

Known issues

Regions of bad pixels in Detector 1 and time varying quantum efficiency in detector 16 renders these detectors the most problematic. The regions of poorer noise statistics are evident in the confidence images. We are aware of a possible offset in the photometric calibration of the Y-band data (of the order of 0.1 mag too bright) and we expect this to be improved upon in subsequent data releases.

For the XMM1 tiles the central position was shifted slightly between early observations and the later ones. The tile centre is given by the TL_RA and TL_DEC keywords in the header information and these provide the centre of the given tile.

Previous Releases

The first public data release of the VIDEO programme via the ESO Science Archive Facility, the VIDEO-XMM Data Release Number 1, Date: 25.07.2011, contained 97 individual (i.e. single-OB) tile images in the VIDEO-XMM3 field and their associated single-band source lists. This release (VIDEO-XMM DR2) adds 156 single-OB images in the VIDEO-XMM fields 1, 2 and 3. DR2 is the first release that includes deep co-added images of the XMM3 field including the associated single-band source lists and the resulting deep photometric catalogue where the ZYJHKs data have been merged.

Data Format

Files Types

Here we release deep stacked images and associated aperture matched catalogues. The naming convention for these is:

```
video_er1_${POSITION}_mosaic_${FILTERNAME}_deepimage_${MFID}.fits.fz  
video_er1_${POSITION}_mosaic_${FILTERNAME}_deepconf_${MFID}.fits.fz  
video_er1_${POSITION}_mosaic_${FILTERNAME}_jpeg_${MFID}.jpg  
video_er1_${POSITION}_mosaic_${FILTERNAME}_cat_${MFID}.fits
```

where `POSITION` is a string like 02h26-004d43 for the XMM3 field.

`FILTERNAME` is z,y,j,h,ks AND `MFID` is the multiframeID in our database.

We note that the jpg images are provided for a quick-look at both the science frames and the confidence images.

The single-tile images, confidence maps and catalogues have the following naming convention:

```
v${DATE}_${OBS_NUMBER}_st_tl.fits.fz  
v${DATE}_${OBS_NUMBER}_st_tl_conf.fits.fz  
v${DATE}_${OBS_NUMBER}_st_tl_cat.fits
```

for the tiled image, the confidence image and the source list respectively. `OBS_NUMBER` refers to the final observation number for the individual paw-prints which contribute to the tile.

Catalogue Columns

A complete list of catalogue columns for the single-band deep-stack source catalogues is given in Table 3 and Table 4 gives the columns for the Ks-band-selected band-merged catalogue. Table 5 gives the column names of the source catalogues derived from individual single tiles derived using the CASU source extraction pipeline (see <http://casu.ast.cam.ac.uk/surveys-projects/vista/technical/catalogue-generation>).

A brief description of these columns now follows.

Source IDs (ObjID, IAUNAME, SOURCEID)

ObjID gives the unique object identifier for the object in the given filter for this DR1. These identifiers are subject to change in future releases.

IAUNAME is the unique source identifier following the IAU convention, and comprising the survey identifier (“VID”) and then the (truncated) RA and Dec of each object. SOURCEID is the unique source number for the merged catalogue.

Positional Parameters (X, Y, RA, Dec)

X and Y report the position of the object in pixels on the relevant image (which applies to all the stacks, as they share the same coordinate system). RA and Dec (and RA200 and DEC2000 in the merged catalogue) in equinox J2000 are based on the 2MASS astrometric system as described in Jarvis et al. (2013).

Magnitude measurements and errors

Aperture magnitudes are measured in 2” and 5.7” fixed diameter apertures (APERMAGNOAPERCORR3 and APERMAGNOAPERCORR6 columns). We also supply seeing corrected “real” aperture magnitudes for these apertures (APERMAG3 and APERMAG6 columns) that provide the photometry in a seeing convolved 2 and 5.7” aperture. The uncertainties on aperture magnitudes are given in APERMAG3ERR, APERMAG6ERR, APERMAGNOAPERCORR3 and APERMAGNOAPERCORR6. The Petrosian magnitude and the associated error are also provided in columns denoted PETROMAG and PETROMAGERR respectively. SExtractor MAG_AUTO magnitudes and uncertainties are denoted as AUTOMAG and AUTOMAGERR.

To correct for correlated noise in the images, the errors for fixed aperture photometry are estimated by measuring the RMS flux in randomly placed apertures of the same size, then adding Poisson errors based on object counts. Errors for Petrosian fluxes are corrected by scaling the random (non-Poisson) component of the error (as estimated by SExtractor); the scaling factor is estimated by comparing the median SExtractor error for randomly placed 5 arcsec apertures with the actual standard deviation of fluxes in those apertures.

Object flags and source classifier (ERRBITS, CLASSTAT, MERGEDCLASS, HALFRAD, EBV)

The SExtractor flag is provided in the column denoted ERRBITS, indicating if an object is blended or has otherwise corrupted measurements. The best possible object sample are those objects which have ERRBITS = 0.

CLASSTAT gives the SExtractor star/galaxy classification statistic, where 0-galaxy and 1-star. The MERGEDCLASS column in the band-merged catalogue is the weighted average of the individual CLASSTAT parameters for that object.

HALFRAD is the SExtractor measurement of the half-light radius in the given filter.

The galactic reddening E(B-V) is reported at the coordinates of each object in the merged catalogue, computed using the Schlegel et al. (1998) dust maps.

In Jarvis et al. objects were discarded when measuring the completeness and reliability of the catalogues using the HALOFLAG term. This flag does not appear in the released catalogues as such a keyword is subject to interpretation and definition. We encourage potential users of the data to define their own object masks according to their science aims.

Table 3. Complete list of supplied catalogue columns in the deep-stacked single-band catalogues.

| Column Number | Column Name | Description |
|---------------|-------------|-----------------------------------|
| 1 | OBJID | Unique Source Identifier |
| 2 | FILTERID | Filter (1=Z; 2=Y; 3=J; 4=H; 5=Ks) |
| 3 | X | X coordinate of detection |

| | | |
|----|-----------------------|---|
| 4 | Y | Y coordinate of detection |
| 5 | RA | Right Ascension in Decimal Degrees |
| 6 | DEC | Declination in Decimal Degrees |
| 7 | APERMAGNOAPERCORR3 | Fixed Aperture magnitude (2" diameter", AB) |
| 8 | APERMAGNOAPERCORR3ERR | Error in calibrated aperture magnitude 3 (2" diameter", AB) |
| 9 | APERMAG3 | Aperture magnitude corrected to 2" diameter, AB |
| 10 | APERMAG3ERR | Error on Aperture magnitude (2", diameter AB) |
| 11 | APERMAGNOAPERCORR6 | Fixed Aperture magnitude (5.7" diameter, AB) |
| 12 | APERMAGNOAPERCORR6ERR | Error in calibrated aperture magnitude 6 (5.7" diameter", AB) |
| 13 | APERMAG6 | Aperture magnitude corrected to 5.7" diameter, AB |
| 14 | APERMAG6ERR | Error on Aperture magnitude (5.7" diameter), AB |
| 15 | AUTOMAG | Extended source magnitude (SExtractor MAG_AUTO) |
| 16 | AUTOMAGERR | Extended source mag error (SExtractor MAG_AUTO) |
| 17 | HALFRAD | Sextractor Half-light radius |
| 18 | PETROMAG | Petrosian Magnitude (AB) |
| 19 | PETROMAGERR | Error on Petrosian magnitude |
| 20 | ERRBITS | Sextractor Flag |
| 21 | CLASSTAT | Sextractor Star/Galaxy classification (0-galaxy, 1-star) |

Table 4. Complete list of supplied catalogue columns in the merged table "VID-EO_XMM3_Kselected_merged_DR1.fits"

| Column Number | Column Name | Description |
|---------------|--------------|---|
| 1 | IAUNAME | IAU-formatted name, prefixed with "VID" |
| 2 | SOURCEID | Unique source identifier |
| 3 | RA2000 | Right Ascension in Decimal Degrees |
| 4 | DEC2000 | Declination in Decimal Degrees |
| 5 | MERGEDCLASS | The average of the CLASSTAT statistic across all bands weighted by $(1/mag_error)^2$ |
| 6 | EBV | Galactic reddening E(B-V) based on Schegel et al. (1998) dust maps |
| 7 | ZMYPNT | Point source colour Z-Y (using aperMag3) |
| 8 | ZMYPNTERR | Error on point source colour Z-Y |
| 9 | ZMYEXT | Extended source colour Z-Y (using aperMagNoAperCorr3) |
| 10 | ZMYEXTERR | Error on extended source colour Z-Y |
| 11 | YMJPNT | Point source colour Y-J (using aperMag3) |
| 12 | YMJPNTERR | Error on point source colour Y-J |
| 13 | YMJEXT | Extended source colour Y-J (using aperMagNoAperCorr3) |
| 14 | YMJEXTERR | Error on extended source colour Y-J |
| 15 | JMHPNT | Point source colour J-H (using aperMag3) |
| 16 | JMHPNTERR | Error on point source colour J-H |
| 17 | JMHEXT | Extended source colour J-H (using aperMagNoAperCorr3) |
| 18 | JMHEXTERR | Error on extended source colour J-H |
| 19 | HMKSPNT | Point source colour H-Ks (using aperMag3) |
| 20 | HMKSPNTERR | Error on point source colour H-Ks |
| 21 | HMKSEXT | Extended source colour H-Ks (using aperMagNoAperCorr3) |
| 22 | HMKSEXTERR | Error on extended source colour H-Ks |
| 23 | ZPETROMAG | Z Petrosian magnitude (AB) |
| 24 | ZPETROMAGERR | Z Petrosian magnitude error (AB) |

| | | |
|----|------------------------|--|
| 25 | ZAPERMAG3 | Z Aperture magnitude corrected to 2" diameter, AB |
| 26 | ZAPERMAG3ERR | Z Error on Aperture magnitude (2", diameter AB) |
| 27 | ZAPERMAG6 | Z Aperture magnitude corrected to 5.7" diameter, AB |
| 28 | ZAPERMAG6ERR | Z Error on Aperture magnitude (5.7" diameter), AB |
| 29 | ZAPERMAGNOAPERCORR3 | Z Fixed Aperture magnitude (2" diameter", AB) |
| 30 | ZAPERMAGNOAPERCORR3ERR | Error in default extended source Z aperture mag, 2.0 arcsec aperture |
| 31 | ZAPERMAGNOAPERCORR6 | Z Fixed Aperture magnitude (5.7' diameter, AB) |
| 32 | ZAPERMAGNOAPERCORR6ERR | Error in extended source Z aperture mag (5.7 arcsec aperture) |
| 33 | ZERRBITS | Z SExtractor Flag |
| 34 | ZCLASSSTAT | SExtractor Star/Galaxy classification (0-galaxy, 1-star) |
| 35 | ZAUTOMAG | Extended source Z mag (SExtractor MAG_AUTO) |
| 36 | ZAUTOMAGERR | Extended source Z mag error (SExtractor MAG_AUTO) |
| 37 | ZHALFRAD | SExtractor half-light radius in Z band |
| 38 | YPETROMAG | Y Petrosian magnitude (AB) |
| 39 | YPETROMAGERR | Y Petrosian magnitude error (AB) |
| 40 | YAPERMAG3 | Y Aperture magnitude corrected to 2" diameter, AB |
| 41 | YAPERMAG3ERR | Y Error on Aperture magnitude (2", diameter AB) |
| 42 | YAPERMAG6 | Y Aperture magnitude corrected to 5.7" diameter, AB |
| 43 | YAPERMAG6ERR | Y Error on Aperture magnitude (5.7" diameter), AB |
| 44 | YAPERMAGNOAPERCORR3 | Y Fixed Aperture magnitude (2" diameter", AB) |
| 45 | YAPERMAGNOAPERCORR3ERR | Error in default extended source y aperture mag, 2.0 arcsec aperture |
| 46 | YAPERMAGNOAPERCORR6 | Y Fixed Aperture magnitude (5.7" diameter, AB) |
| 47 | YAPERMAGNOAPERCORR6ERR | Error in extended source y aperture mag (5.7 arcsec aperture) |
| 48 | YERRBITS | Y SExtractor Flag |
| 49 | YCLASSSTAT | Y SExtractor Star/Galaxy classification (0-galaxy, 1-star) |
| 50 | YAUTOMAG | Extended source Y mag (SExtractor MAG_AUTO) |
| 51 | YAUTOMAGERR | Extended source Y mag error (SExtractor MAG_AUTO) |
| 52 | YHALFRAD | SExtractor half-light radius in Y band |
| 53 | JPETROMAG | J Petrosian magnitude (AB) |
| 54 | JPETROMAGERR | J Petrosian magnitude error (AB) |
| 55 | JAPERMAG3 | J Aperture magnitude corrected to 2" diameter, AB |
| 56 | JAPERMAG3ERR | J Error on Aperture magnitude (2", diameter AB) |
| 57 | JAPERMAG6 | J Aperture magnitude corrected to 5.7" diameter, AB |
| 58 | JAPERMAG6ERR | J Error on Aperture magnitude (5.7" diameter), AB |
| 59 | JAPERMAGNOAPERCORR3 | J Fixed Aperture magnitude (2" diameter", AB) |
| 60 | JAPERMAGNOAPERCORR3ERR | Error in default extended source j aperture mag, 2.0 arcsec aperture |
| 61 | JAPERMAGNOAPERCORR6 | J Fixed Aperture magnitude (5.7 diameter, AB) |
| 62 | JAPERMAGNOAPERCORR6ERR | Error in extended source j aperture mag (5.7 arcsec aperture) |

| | | |
|----|-------------------------|--|
| 63 | JERRBITS | J SExtractor Flag |
| 64 | JCLASSSTAT | J SExtractor Star/Galaxy classification (0-galaxy, 1-star) |
| 65 | JAUTOMAG | Extended source J mag (SExtractor MAG_AUTO) |
| 66 | JAUTOMAGERR | Extended source J mag error (SExtractor MAG_AUTO) |
| 67 | JHALFRAD | SExtractor half-light radius in J band |
| 68 | HPETROMAG | H Petrosian magnitude (AB) |
| 69 | HPETROMAGERR | H Petrosian magnitude error (AB) |
| 70 | HAPERMAG3 | H Aperture magnitude corrected to 2" diameter, AB |
| 71 | HAPERMAG3ERR | H Error on Aperture magnitude (2", diameter AB) |
| 72 | HAPERMAG6 | H Aperture magnitude corrected to 5.7" diameter, AB |
| 73 | HAPERMAG6ERR | H Error on Aperture magnitude (5.7" diameter), AB |
| 74 | HAPERMAGNOAPERCORR3 | H Fixed Aperture magnitude (2" diameter, AB) |
| 75 | HAPERMAGNOAPERCORR3ERR | Error in default extended source h aperture mag, 2.0 arcsec aperture |
| 76 | HAPERMAGNOAPERCORR6 | H Fixed Aperture magnitude (5.7" diameter, AB) |
| 77 | HAPERMAGNOAPERCORR6ERR | Error in extended source h aperture mag (5.7 arcsec aperture) |
| 78 | HERRBITS | H SExtractor Flag |
| 79 | HCLASSSTAT | H SExtractor Star/Galaxy classification (0-galaxy, 1-star) |
| 80 | HAUTOMAG | Extended source H mag (SExtractor MAG_AUTO) |
| 81 | HAUTOMAGERR | Extended source H mag error (SExtractor MAG_AUTO) |
| 82 | HHALFRAD | SExtractor half-light radius in H band |
| 83 | KSPETROMAG | Ks Petrosian magnitude (AB) |
| 84 | KSPETROMAGERR | Ks Petrosian magnitude error (AB) |
| 85 | KSAPERMAG3 | Ks Aperture magnitude corrected to 2" diameter, AB |
| 86 | KSAPERMAG3ERR | Ks Error on Aperture magnitude (2", diameter AB) |
| 87 | KSAPERMAG6 | Ks Aperture magnitude corrected to 5.7 diameter, AB |
| 88 | KSAPERMAG6ERR | Ks Error on Aperture magnitude (5.7" diameter), AB |
| 89 | KSAPERMAGNOAPERCORR3 | Ks Fixed Aperture magnitude (2" diameter", AB) |
| 90 | KSAPERMAGNOAPERCORR3ERR | Error in default extended source ks aperture mag, 2.0 arcsec |
| 91 | KSAPERMAGNOAPERCORR6 | Ks Fixed Aperture magnitude (5.7" diameter", AB) |
| 92 | KSAPERMAGNOAPERCORR6ERR | Error in extended source ks aperture mag (5.7 arcsec aperture) |
| 93 | KSERRBITS | Ks SExtractor Flag |
| 94 | KSCLASSSTAT | Ks SExtractor Star/Galaxy classification (0-galaxy, 1-star) |
| 95 | KAUTOMAG | Extended source Ks mag (SExtractor MAG_AUTO) |
| 96 | KAUTOMAGERR | Extended source Ks mag error (SExtractor MAG_AUTO) |
| 97 | KSHALFRAD | SExtractor half-light radius in Ks band |

The individual-tile catalogue format was derived from similar APM/SuperCOSMOS/INTWFC/CIRSI analysis which produced 32 4-byte parameters per detected object. This was first enhanced for WFCAM to an 80 4-byte parameter set to include extra parameters for flux estimation and error estimates, and this has now been further refined for the

VIRCAM catalogues. The following tables cover the VIRCAM standard and further processing pipeline output catalogues, where for simplicity all derived parameters are stored as floating point numbers even though some of them are more naturally integers. See <http://casu.ast.cam.ac.uk/surveys-projects/vista/technical/catalogue-generation> for more information.

Table 5. Complete list of supplied catalogue columns for the sources lists released with the individual tiles.

| No | Name | Column Description |
|-------|--|---|
| 1 | Seq No. | Running number for ease of reference, in strict order of image detections |
| 2 | Isophotal flux | Standard definition of summed flux within detection isophote, apart from detection filter is used to define pixel connectivity and hence which pixels to include. This helps to reduce edge effects for all isophotally derived parameters. |
| 3 | X coord | Intensity-weighted isophotal centre-of-gravity in X |
| 4 | Error in X | Estimate of centroid error |
| 5 | Y coord | Intensity-weighted isophotal centre-of-gravity in Y |
| 6 | Error in Y | Estimate of centroid error |
| 7 | Gaussian sigma | These are derived from the three general intensity-weighted second moments |
| 8 | Ellipticity | The equivalence between them and a generalised elliptical Gaussian |
| 9 | Position angle | Position angle of the isophote |
| 10-17 | Areal profile 1-8 | Number of pixels above a series of threshold levels relative to local sky. Levels are set at T, 2T, 4T, 8T . . . 128T where T is the threshold. These can be thought of as a radial profile. Note that for now, deblended, i.e. overlapping images, only the first areal profile is computed and the rest are set to -1 flagging the difficulty of computing accurate profiles for blended images this parameter is used to flag the start of the sequence of the deblended components by setting the first in the sequence to 0. |
| 18 | Peak height | In counts relative to local value of sky - also zeroth order aperture flux |
| 19 | Error in pkht | Error in Peak Height |
| 20-45 | Aperture flux 1-13 and error on aperture flux 1-13 | These are a series of different radii soft-edged apertures designed to adequately sample the curve-of-growth of the majority of images and to provide fixed-sized aperture fluxes for all images. The scale size for these apertures is selected by defining a scale radius $\langle FWHM \rangle$ for site+instrument. In the case of VIRCAM this "core" radius (r_{core}) has been fixed at 1.0 arcsec for convenience in inter-comparison with other datasets. A 1.0 arcsec radius is equivalent to 3.0 pixels for normal data. In 0.8 arcsec seeing an r_{core} -radius aperture contains roughly 75% of the total flux of stellar images. The aperture fluxes are sky-corrected integrals (summations) with a soft-edge (i.e. pro-rata flux division for boundary pixels). However, for overlapping images they are more subtle than this since they are in practice simultaneously fitted top-hat functions, to minimise the effects of crowding. Images external to the blend are also flagged and not included in the large radius summations. Starting with parameter 20 the radii are: $1/2 \times r_{core}$, $1/\sqrt{2} \times r_{core}$, r_{core} , $\sqrt{2} \times r_{core}$, $2 \times r_{core}$, $2\sqrt{2} \times r_{core}$, $4 \times r_{core}$, $5 \times r_{core}$, $6 \times r_{core}$, $7 \times r_{core}$, $8 \times r_{core}$, $10 \times r_{core}$, $12 \times r_{core}$ We recommend using Aperture flux 3 if a single number is required to represent the flux for ALL images - this aperture has a radius of $r_{core} = 1$ arcsec. |

| | | |
|----|------------------|---|
| 46 | Petrosian radius | Petrosian radius as defined in Yasuda et al. 2001 AJ 112 1104 |
| 47 | Kron radius | Kron radius as defined in Bertin and Arnouts 1996 A&A Supp 117 393 |
| 48 | Hall radius | Hall image scale radius eg. Hall & Mackay 1984 MNRAS 210 979 |
| 49 | Petrosian flux | Petrosian flux within circular aperture to $k \times r_p$ with $k=2$ |
| 50 | Error in flux | Error on Petrosian flux |
| 51 | Kron flux | Flux within circular aperture to $k \times r_k$ with $k=2$ |
| 52 | Error in flux | Error on Kron flux |
| 53 | Hall flux | Flux within circular aperture to $k \times r_h$ with $k=5$; alternative total flux |
| 54 | Error in flux | Error on Hall flux |
| 55 | Error bit flag | Bit pattern listing various processing error flags initially set to the no. of bad pixels within aperture of radius "rcore"- note this can be fractional due to soft-edged apertures |
| 56 | Sky level | Local interpolated sky level from background tracker |
| 57 | Sky rms | Local estimate of variation in sky level around image |
| 58 | Av conf | Average confidence level within default rcore aperture useful for spotting spurious outliers in various parameter selection spaces |
| | | The following are accreted after standard catalog generation |
| 59 | RA | Right Ascension in radians (J2000) |
| 60 | Dec | Declination in radians(J2000) |
| 61 | Classification | Flag indicating most probable morphological classification: eg. -1 stellar, +1 non-stellar, 0 noise, -2 borderline stellar (Saturated images can be flagged by comparing the peak height + local sky with the SATURATE keyword in the header.) |
| 62 | Statistic | An equivalent $N(0,1)$ measure of how stellar-like an image is, used in deriving parameter 61 in a 'necessary but not sufficient' sense. Derived mainly from the curve-of-growth of flux using the well-defined stellar locus as a function of magnitude as a benchmark (see Irwin et al. 1994 SPIE 5493 411 for more details). |
| 63 | MJDoff | The number of minutes to be added to MJD_DAY in order to get the MJD of a given object. Hence for the i th object: $MJD_i = MJD_DAY + MJDoff_i / 1440.0$ |
| | | |
| 64 | Blank | Blank |
| 65 | Blank | Blank |
| 66 | Blank | Blank |
| 67 | Blank | Blank |
| 68 | Blank | Blank |
| 69 | Blank | Blank |
| 70 | Blank | Blank |
| 71 | Blank | Blank |
| 72 | Blank | Blank |
| 73 | Blank | Blank |
| 74 | Blank | Blank |
| 75 | Blank | Blank |
| 76 | Blank | Blank |
| 77 | Blank | Blank |
| 78 | Blank | Blank |
| 79 | Blank | Blank |
| 80 | Blank | Blank |

Acknowledgements

The VIDEO Survey is described in Jarvis et al. 2013, MNRAS, 428, 1281 and any publications using the VIDEO data should reference this article.

Please use the following statement in your articles when using these data:

Based on data products from observations made with ESO Telescopes at the La Silla Paranal Observatory under programme ID 179.A-2006.

References

- Bertin E. & Arnouts S., 1996, ApJS, 117, 393
Irwin M.J., et al., 2004, SPIE, 5493, 411
Jarvis M.J., et al., 2013, MNRAS, 428, 1281
Schlegel D.J., Finkbeiner D.P. & Davis M., 1998, ApJ, 500, 525
Skrutskie M.F., et al., 2006, AJ, 131, 1163