provided by T. Shanks, submitted on

ഗ 0 ഗ ш

ESO Phase 3 Data Release Description

Data Collection ATLAS DR5

Release Number 5

T Shanks **Data Provider** <22.10.2025> **Date**

Authors: T Shanks, N Metcalfe

Physics Department, University of Durham, South Road, Durham, DH1 3LE, UK

C Gonzalez-Fernandez, E Gonzalez-Solares, MI Irwin

CASU, Institute of Astronomy, Univ. of Cambridge, Madingley Road, Cambridge, CB3 0HA, UK NJG Cross, MA Read

WFAU, Institute for Astronomy, Univ. of Edinburgh, Blackford Hill, Edinburgh EH9 3HJ, UK

Abstract

The new data released here are the final VLT Survey Telescope (VST) ATLAS reduced +stacked images and associated source lists taken from observations by VST from October 2017 to the end of the survey in 🔁 January 2021 under ESO id 177.A-3011. These new data mainly comprise the ATLAS NGC extension at Dec<-20deg. Since DR5 is the final ATLAS data release, this document also summarises the full ATLAS surby vey content and characteristics, covering $\sim 4700 \text{deg}^2$ of sky in the SGC and NGC in the SDSS u, g, r, i, z bands and reaching similar depth ($r\sim22.5$) as the SDSS in the Northern Hemisphere. Basic data reduction was done at the Cambridge Astronomical Surveys Unit (CASU). Median FWHM seeing ranges from 0.8arcsec 5 in i,z to 1.1 arcsec in u. Each ATLAS tile comprises a stacked pawprint composed of 2 pawprints offset by 25 arcsec in X and 85 arcsec in Y which takes out most of the inter-chip gaps between the 32 OmegaCAM 25 arcsec in X and 85 arcsec in Y which takes out most of the little composition of the little composition. The CCDs. There are also 2 arcmin overlaps in both RA and Dec between tiles to allow cross-calibration. The ATLAS survey is particularly aimed at survey cosmology but can be exploited for many other branches of E extragalactic and Galactic astronomy. Its wide wavelength coverage from the *u* to the *z* bands complements the WETA Hard of the coverage from the *u* to the *z* bands complements the WETA Hard of the coverage from the *u* to the *z* bands complements the WETA Hard of the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands complements the coverage from the *u* to the *z* bands coverage from the *u* to the *u* to the *z* bands coverage from the *u* to the *z* band ments the VISTA Hemisphere and VIKING Surveys in the YJHK bands. A description of the ATLAS survey is given by Shanks et al (2015, MNRAS, 451, 4238). DR5 presents new and updated CASU images and source given by Shanks et al (2015, MNRAS, 451, 4238). DR5 presents new and updated CASU images and sourc lists and also a full re-release of multi-band catalogue data from the Wide Field Astronomy Unit (WFAU) at Edinburgh, representing our "best" data. Any other un-updated data is still available within DR4 and earlier releases. DR5 data also includes the second of the earlier releases. DR5 data also includes the final ATLAS global photometry calibration.

 $^{ t Q}$ Table 1 summarises the new contents and basic characteristics of the ATLAS DR5 data release:

Table 1 – Basic VST ATLAS DR5 Summary									
	u	u	g	r	i	z			
Exposure (s)	2x60s	2x120s	2x50s	2x45s	2x45s	2x45s			
No. of Tiles	34 (1)	797 (797)	812 (798)	808 (797)	77 (1)	69 (1)			
No. of Stacked	34	797	812	808	77	69			
Pawprints									
~Area (deg²)	32	745	759	755	72	64			
Median Mag	21.70	22.39	23.14	22.65	21.83	20.65			
Lim.									
Median Sky	22.38	22.41	22.16	21.19	19.69	18.77			
Bri.									
20mag e ⁻ /s	29	29	177	160	101	29			
Median See-	1.07	1.04	0.94	0.98	0.86	0.99			
ing (")									

Notes: Rows (2,3:, The first number is tiles contained in DR5. In brackets is the number new to DR5. In DR5 the number of stacked pawprints is now the same as the number of tiles covered since only the best observations are included. Row (5): Median 5σ AB point source magnitude limit in 2 arcsec aperture for ATLAS DR5. Row (6): Median sky brightness in AB mags/arscec². Row (7): fluxes for AB 20 mag point sources, normalized to airmass 1.3. Median seeing (FWHM in arcsec). The magnitude limit, sky brightness and seeing are measured from chip #12, which is one of the four closest to the centre of the detector. All magnitude limits and sky brightnesses are in a system close to SDSS AB.

Table 2 summarises the final, overall, characteristics of the full ATLAS survey:

Table 2 – Final Overall VST ATLAS Summary									
	u	u	g	r	i	Z			
Exposure (s)	2x60s	2x120s	2x50s	2x45s	2x45s	2x45s			
No. of Tiles/	4276	797	5073	5073	5073	5073			
Stacked Pawprints									
~Area (deg²)	3994	745	4739	4739	4739	4739			
Median Mag Lim.	21.95	22.39	23.22	22.70	22.05	20.84			
Median Sky Bri.	22.37	22.41	22.01	20.98	19.81	18.82			
20mag e ⁻ /s	29	29	177	160	101	29			
Median Seeing (")	1.02	1.04	0.93	0.89	0.79	0.82			

Notes: Row (1): The u exposure of 2x60s applies to all tiles except those in the NGC with Dec<-20° where the u exposure was 2x120. Rows (2,3): These are the total number of tiles and their area contained in the full AT-LAS survey. Row (4): Median 5σ AB point source magnitude limit in 2 arcsec aperture for full survey. Row (5): Median sky brightness in AB mags/arscec².for full survey. Row (6): fluxes for AB 20 mag point sources, normalized to airmass 1.3. Row (7): Median seeing (FWHM in arcsec) for full survey. The magnitude limit, sky brightness and seeing are measured from chip #12, which is one of the four closest to the centre of the detector.All magnitude limits and sky brightnesses are in a system close to SDSS AB.

These exposure times are generally longer than the $\sim 55 \mathrm{s}$ of SDSS to take into account the 0.21 arcsec pixels of VST OmegaCam compared to the 0.4 arcsec SDSS pixels and slightly brighter skies for ATLAS in i and z to produce approximately similar S/N as SDSS in all bands. Note that ATLAS OBs are generally composed of a concatenation of 17 tiles in a given band, taken in fixed Declination strips in the direction of increasing RA. The ugr images are taken in dark time and the iz images are taken in grey/bright time.

Images and source lists for all ESO Grades are being supplied. The specified survey quality is limited to ESO Grade A, B and occasionally C, in cases where only a single tile in a 17 tile concatenation was at C grade. Otherwise C and D grade tiles have been repeated. Note that $\sim 4\%$ of images have no ESO grades available. As for DR4, DR5 only contains the best stacked pawprints and this means the total number of DR5 stacked pawprints is the same as the total number of tiles observed (see also Table 1). The TL_RA and TL_DEC header keywords are identical for all data belonging to the same tile.

DR5 provides ugr coverage below a declination of -20 deg (between RA 10h and 16h), which was absent from DR4 (an area of 797 tiles). The iz data in this area were released in DR4. The u data in this region has double the exposure of the rest of the survey (2x120s), improving the u limit by \sim 0.5 mag. In the rest of the ATLAS area, a similar doubling of exposure time to 4x60s can be obtained by including the 2x60s u exposures of the ATLAS Chilean Extension (ACE) of F. Barrientos et al. (in prep). Note this survey is not an ESO Public Survey. DR5 also provides some improved observations over the rest of the survey area.

DR4 and DR5 combined now provide complete ugriz coverage over the whole $\sim 4700~deg^2$ ATLAS area. ncluding 2597 stacked pawprints + 2597 confidence maps + 2597 source lists, the CASU component of DR5 comprises 7791 files, occupying a total of ~ 0.68 Tb. The complete survey is composed of 27962 pawprints, confidence maps and source lists, making a total of 83886 files. The total survey size is ~ 6.9 Tb.

Fig. 1 shows the DR5 ATLAS sky coverage by filter and Fig. 2 shows the *ugriz* sky coverage for the full ATLAS survey.





Figure 1: ugriz sky coverage of ATLAS DR5.



Figure~2: Final~ugriz~sky~coverage~offull~ATLAS~survey~including~DR5.




Fig ure 3: Seeing (FWHM) distribution for the tiles released in DR5 (taken from chip #12). The u band is split into 2x120s (blue) and 2x60s (cyan).




Figure 4. As Fig. 3 but for the entire ATLAS survey (DR4+DR5).

Release Content

The CASU imaging data comprises the combination of the two individual pawprint images which goes to make an ATLAS stacked pawprint in each tile. Each file is in a multi-extension fits (MEF) format with an extension for each of the 32 OmegaCam CCDs in the stacked pawprint. Individual CCDs originally contained 2048x4096 pixels and the stacked pawprint extensions contain approximately 2165x4500 pixels to cover the two 25"x85" offset CCDs that make up each extension in a stack. Along with the imaging data, we are also releasing statistical confidence maps in the same format. The seeing is specified to be <1.4 arcsec FWHM and the distributions by passband are shown in Fig. 3 for DR5 and in Fig. 4 for the entire survey. The distribution of limiting magnitudes at the 5σ detection level by passband is shown in Fig. 5 for DR5, and in Fig. 6 for the entire survey. The total area covered by DR5 is ~760 deg² in ugr (with ~745 deg² being the new data below dec -20 deg) and ~70 deg² in iz (mostly repeat observations). The 2-pointing dither leaves 14 small (2x80x20arcsec²) holes amounting to 1/3% of the total area.

Also included in DR5 are bandmerged catalogue data from WFAU (see Data Format section below). Note that the WFAU DR5 catalogue data are a complete re-release of the whole ATLAS survey and so supersede WFAU DR4. The DR5 release consists of 5074 files occupying ~115Gb.





Figure 5: ATLAS (Global Calibration) 5σ AB magnitude limit distributions for DR5 point sources in u (2x120s – blue) (2x60s - cyan), g (green), r (red), i (purple) and z (grey). Data are taken from chip #12. The median magnitude limits are given in Table 1.

The CASU source list data covers the same pixel areas as the stacked pawprints (see above). The derived object source lists are also stored as multi-extension FITS files using FITS binary tables, one for each image extension with the primary header unit containing the telescope and observation-specific information. The source list extension headers contain a copy of the relevant detector-specific information. Each detected object has an attached set of descriptors, forming the columns of the binary table and summarising derived position, shape and intensity information. During further processing stages ancillary information such as the sky properties, seeing and so on are derived from the source lists and stored in the FITS headers attached to each source list extension. All derived parameters are stored as floating point numbers. A full description of the source list columns is given at

http://apm49.ast.cam.ac.uk/surveys-projects/vst/technical/catalogue-generation



Release Notes

Astrometric calibration is via the numerous unsaturated 2MASS point sources available in each tile. By stacking residuals from a series of standard Tangent Plane astrometric fits based on 2MASS we can see (as in the example in Fig. 4 below) that there are no significant astrometric distortions over the whole field of view. The individual detector astrometric solutions achieve rms accuracies of around 70-80mas per star - generally dominated by rms errors in 2MASS stars. Even at high Galactic latitudes there are sufficient calibrators to give systematic residuals at the ~25mas level per detector. The global systematics from stacking multiple solutions are better than this as can be seen in Fig. 4. A Tangent Plane projection (TAN) is being used for all data products.




Figure 7. Astrometric VST-2MASS residuals for stars as a function of distance from tile centre.

The original ATLAS(ESO) photometric calibration of each pointing is based on the limited number of standard fields observed by VST each night. This calibration was in a VST Vega-like system, but, as the average standard star SED and the detector response drops rapidly in the UV, it would be surprising if there were not issues in at least the u-band calibration. Note that these zero-points are still based on the original source lists rather than the illumination-correction fixed source lists (see below). An improved AB calibration for DR2+DR3 was based on the AAVSO Photometric All-Sky Survey (APASS-http://www.aav-so.org/apass) is described in the Data Reduction and Calibration section below where the further improved calibration for DR4 and DR5 based on Gaia photometry in griz and tile overlaps in u is also de-

scribed.

Data Reduction and Calibration

The data processing of the science products released in this data release mainly comes from version 1.0 of the VST Data Flow System (VDFS) pipeline running at CASU, identified with keyword PROCSOFT=`omegacam version 1.0'. Unlike VPHAS, there was no use of the nebulosity filter prior to source extraction. The reflection artefacts around heavily saturated bright stars can generate spurious detections but the contamination is reduced if source lists are matched during band-merging. CASU source lists for individual pawprints are supplied with no attempt to reject overlaps (but note that overlaps are addressed in the WFAU bandmerged catalogues - see below). The astrometric reference catalogue is 2MASS. The image illumination corrections are made from a comparison of each month of data from all VST public surveys with the APASS survey. This correction has been applied to source list photometry but not to image pixels. CASU have developed software to apply the illumination correction to image pixels. Since scattered light is also present in dark sky science images the optimum way to use this correction depends on the end-user requirements so this correction is not routinely applied to the stored images). Stellar aperture corrections are supplied for each photometric aperture used in the source list and these can be used to estimate total fluxes or magnitudes for stars. PSF magnitudes will be produced for the final data release. Source fluxes in the binary tables are in ADUs and require corrections for aperture loss, airmass (relative to unity), and application of the appropriate magnitude zeropoint. The relevant information is supplied in the source list headers. No correction for Galactic extinction has been applied or supplied, in part because the correction is specific to the extinction model adopted.




Figure 8. The illumination correction in the i-band as measured by residuals from APASS photometry. The released source lists have had the illumination correction applied based on source position in the image plane.

Data Quality

The astrometric calibration is uniform across the survey to an rms accuracy of 25mas relative to 2MASS. Indeed, various fields scattered across the ATLAS area have already been used as the basis for 2dF fibre observations with no astrometric problem. The main problem in the released images is a non-uniform photometric response across the field caused by the presence of scattered light in the VST flat-fields, with a pattern across the pawprint which typically looks like that shown in Fig. 5. The scattered light is made up of multiple components having different symmetries and scales causing effects ranging in scale from ten arcsec with x-y rectangular symmetry, e.g. due to scattering off masking strips of CCDs, to large fractions of the field due to radial concentration in the optics and to non-astronomical scattered light entering obliquely in flats. The illumination correction removes the dominant reproducible components of this effect in the source lists leaving the zeropoint across the field uniform to $\sim \pm 0.003$ mag. We note that after the recent VST baffling improvements, and particularly after January 2014, the size of the illumination correction required has dropped by more than a factor of two and in current data the range is approximately +/- 0.05 magnitudes. In addition, one detector, #82, otherwise known as CCD #10 in the MEF extensions, had a gain which intermittently varied by up to 0.5 mag until the video board replacement in June 2012; this may not always be taken out by the flatfield.

Quality of magnitudes have been better checked for point sources than for galaxies. It may be possible to extend depth for galaxies by specific smoothing of image before source detection.

Contamination of the source list by spurious sources is at the <5% level down to the limiting magnitudes estimated in the headers. The source lists are estimated to be ~50% complete at the quoted 5σ limiting magnitudes.

Finally, in Fig. 6 we show for a science example, colour-colour diagrams for g<22.5 stars in one ATLAS tile as recently used to select quasars for the 2dF QSO Dark Energy Survey pilot (Chehade et al, 2016, MNRAS, 459, 1179) at the Anglo-Australian Telescope (AAT).




Figure 9. ugr, gri and riz colour-colour diagrams for stellar objects to $g\sim22.5$ in an ATLAS tile as used to select QSO targets for AAT 2dF. The red contours are from SDSS selected stars for comparison.

Previous (DR2, DR3) Outdated Photometric Calibrations

The original (ATLAS-ESO) magnitude zeropoint calibration and error (MAGZPT and MAGZRR in the header) was good to ± 0.05 mag between tiles across the survey, as estimated from a comparison with APASS m<16 stellar magnitudes (see Fig. 7). This zeropoint calibration applied in the Vega magnitude system across all filters and epochs. In this calibration, zeropoints of colour indices may thus be good to $\sim \pm 0.07$ mag. These ATLAS-ESO zeropoints are still supplied in ATLAS headers in DR4 and DR5.




Figure 10. The ugriz distributions of APASS AB-VST Vega offsets for m<16mag stars from ~240 tiles in the SGP GAMA region.

With DR2 and DR3, two new magnitude zeropoints became available based on APASS stellar photometry

to ~16mag. The APASS zeropoint (APASSZPT in the header) and its error (APASSZRR) is based on a comparison of APASS and ATLAS stars in the 13<m<16mag range within each stacked pawprint. The number of stars that contributed to each pawprint zeropoint is given by APASSNUM. The ATLAS-APASS nightly zeropoint and error (NIGHTZPT and NIGHTZRR in the header) is based on the average of all the APASS zeropoints measured on a given night in a particular passband. The number of APASS zeropoints that contributed to this nightly zeropoint is given by NIGHTNUM. These APASS zeropoints are again still supplied in ATLAS headers in DR4 and DR5.

We estimate the accuracy of the ATLAS (APASS) nightly zeropoint from a comparison with SDSS in its $120 deg^2$ overlap area with ATLAS in the NGC. The ATLAS (APASS) - SDSS magnitude standard deviations are ± 0.035 , ± 0.013 , ± 0.013 , ± 0.012 and ± 0.055 mag in ugriz, in most bands a significant improvement over the ATLAS(ESO) - SDSS comparison (± 0.045 , ± 0.027 , ± 0.037 , ± 0.035 and ± 0.073 in ugriz). The ATLAS-APASS nightly zeropoint is therefore recommended for use in ATLAS DR2 and DR3.

For DR4 and DR5, the DR4 Global Calibration is recommended as described below.

DR4/5 Global Calibration

DR4/5 contain the final global calibration for ATLAS photometry. In the *griz* bands, Gaia (Gaia Collaboration et al., 2018, A&A, 616, A1) G, B_p , R_p stellar photometry have been used as the basis of the calibration. To convert to the ATLAS passbands the following colour equations have been used:

$$\begin{split} g_{vst} &= G_{gaia} + 0.35 (Bp-Rp)_{gaia} + 0.3 (Bp-Rp)^2_{gaia} - 0.0085^* (Bp-Rp)^5_{gaia} - 0.08 \text{ [for } 0.2 < (Bp-Rp) < 2.4 \text{]} \\ r_{vst} &= Rp_{gaia} + 0.30 (Bp-Rp)_{gaia} - 0.08 (Bp-Rp)^2_{gaia} + 0.072^* (Bp-Rp)^3_{gaia} + 0.25 \text{ [for } 0.4 < (Bp-Rp) < 2.4 \text{]} \\ i_{vst} &= Rp_{gaia} - 0.03 (Bp-Rp)_{gaia} + 0.03 (Bp-Rp)^2_{gaia} + 0.33 \\ z_{vst} &= Rp_{gaia} - 0.01 (Bp-Rp)_{gaia} - 0.14 (Bp-Rp)^2_{gaia} - 0.022 (Bp-Rp)^3 + 0.30 \end{split}$$

In the u-band we found that the effect of Galactic dust on the Gaia colour terms were too large to apply this method. Therefore we used the ~2arcmin overlaps make a least squares solution via the matrix method of Glazebrook et al (1994, MNRAS, 266, 65). We used overlap stars down to u<17. We assumed ~40% of ATLAS pawprints as anchor fields selected from fields observed on nights that appeared photometric. Initially we used the original ESO photometric zeropoints in the anchor fields but this was ultimately converted from u_Vega to u_AB for the final overall zeropoints by adding +0.86mag to the ESO anchor zeropoints.

The header keywords corresponding to this latest calibration are ATLASZPT and ATLASZRR, representing the new pawprint zeropoint and its error, always in (AB) magnitude units. For ATLASZRR we assume 0.02mag for the griz error and 0.04mag for the u error.

To calculate a stellar magnitude, m_{star} , PHOTZP (based on ATLASZPT) is used as follows:

```
m_{star} = -2.5*log_{10}(APER_FLUX_3) - APCOR3 + PHOTZP
```

where

```
PHOTZP = ATLASZPT - EXTINCT^*(AIRMASS-1) + 2.5*log_{10}(EXPTIME/2).
```

(EXPTIME given in the FITS headers is the sum of the two exposures, but the fluxes are the mean, hence the EXPTIME/2 in the above equation.

PHOTZP is supplied in the headers along with its error PHOTZPER.

In this example, APER_FLUX_3 is the star's flux in aperture 3 (1 arcsec radius) and APCOR3 is the stellar aperture correction appropriate for seeing conditions etc on that pawprint. AIRMASS is usually the average of the airmasses recorded at the start and end of the exposure. EXTINCT is the assumed extinction co-

efficient for that passband. Further details of ATLAS apertures and magnitudes are given by Shanks et al (2015, MNRAS, 451, 4238).

Further tests of the ATLAS photometry calibrations can be found at http://astro.dur.ac.uk/cea/vstatlas/tests/

Known issues

The illumination correction for scattered light in the flatfields has only been applied to source lists and not the image pixels. This problem will have to be addressed before attempting surface photometry of very large galaxies but even then the additive scattered light present in all images may preclude achieving reliable surface photometry at faint magnitudes.

We also note the occasional presence of pickup noise in some observations at the level of ±10ADU caused by guide/wavefront sensor readout happening at same time as science frame read-out. A similar pickup pattern occurs on all 32 detectors and is fixable in post-processing. This problem was (mostly) fixed by some modifications to control software in Autumn 2012.

Note that in observations taken in approximately the first 2 months of the survey, ie before 3/10/11, the 25 arcsec dither in $X(\sim RA)$ between the two pawprints that make up the stacked pawprint was gradually reduced as each of the 17 tiles in a Group for a given RA range was observed, due to a VST technical problem. This means that the main CCD gaps in the Dec direction are not filled in as well as they should be in the stacked pawprint. Data taken after the above date were taken as concatenations and should be unaffected by this problem.

The six DR5 fields observed in January 2021 have one failed OmegaCam CCD chip. Although this exists as a valid FITS extension in the files, it contains no useable data.

For source lists, the correct keyword values that characterise the observation at large (e.g. T/EXPTIME, MJD-OBS, TELESCOP etc) are to be taken from the primary header, and not from the extension header.

Previous Releases

DR1 was the first data release for VST ATLAS. DR1 was superseded by DR2. DR2 therefore improved on the data released in DR1 and included 94 new files for the DR1 period 8/11-9/12 as listed below. The main reason most were omitted was because they were ESO test data. Although "ATLAS depth test" usually appears in the header, this is a misnomer for these files since they were generally made as part of an ESO "Concatenation test" when ATLAS moved to use concatenations of observing blocks. But much of these data is good quality and so they were included in DR2. 6 files that were included in DR1 and are now deprecated/dropped from DR2 because they represent observations whose exposure was interrupted. See the DR2 Release Description for lists of these files.

The DR3 release of CASU imaging and source lists delivered entirely new data covering observations made between 1/10/13 to 31/9/14. These data had therefore to be used alongside the DR2 CASU data, covering the period from survey start to 30/9/14. On the other hand, the DR3 band-merged source lists from WFAU cover all observations taken from the survey start to 30/9/14 and so supersede the ATLAS data released from WFAU in DR2. DR4 was a superseding release that contained only the best observations made up to 31/7/17. DR4 was the first release to contain the new global photometric calibration as described above.

Data Format

File Types

The CASU image files are in multi-extension FITS (MEF) format with an extension for each of the 32 CCDs in each stack. The CASU derived object source lists are also stored in multi-extension FITS files as FITS binary tables, one extension for each image extension with the primary header unit containing the telescope and observation-specific information. The source list extension headers contain a copy of the rele-

vant detector-specific information.

Both CASU images and source list filenames are in the form o20110914_00092 where the night of observation is followed by the ESO VST nightly run number. The suffix_st indicates that the image or source list is based on a stacked pawprint. A suffix _cat indicates that the file corresponds to a source list. A suffix _conf indicates that the file contains the statistical confidence map for the tile. The file type fits.fz indicates Rice compressed FITS file and these can be de-compressed using e.g. CFITSIO funpack.

As with DR4, for DR5 we have only included the list of 'best' files for each tile. Generally, the best files are assumed to be based on the latest observations to take into account repeat observations of the same tile when these had been requested to replace original lower quality C or D grades etc.

As with DR4, we have also fixed a problem with some pawprints caused by the seeing varying significantly between sub-exposures. This caused photometry issues in the inter-chip areas covered by only one sub-exposure in a stacked pawprint resulting in e.g. too many objects being classified as galaxies. This can also cause the seeing in this area of the pawprint to be returned wrongly as 0.0 or -1.0. The affected areas were "grouted" by re-doing star-galaxy classification using parameters from the individual sub-exposure rather than the stack. These stacked pawprints were also checked for issues associated with cosmic ray rejection caused by seeing differences and re-stacked where too many pixels had been falsely rejected.

The DR2, DR3, DR4 and DR5 releases again contain bandmerged source lists produced by the Wide Field Astronomy Unit (WFAU), Edinburgh. The DR5 release contains data for the entire survey, so does not need to be combined with previous releases. These files combine a subset of the parameters as supplied in the CASU lists but now with magnitudes calculated from fluxes and observations merged so that colour indices are provided. Additional parameters include extinction coefficients, merged classification statistics and error bit flagging. The flag, priOrSec allows queries that can select out a seamless catalogue or can be used to identify objects in overlap regions. Also, the PRIMARY_SOURCE column is based on this flag, again identifying the primary object while still including secondary objects. The primary is selected as the source with the best observations or the one closest to the optical axis if all else is equal (see "Source List Columns" below).

The magnitudes are computed based on the ATLAS (Global calibration) zeropoints described above. The data files are based on framesets formed by merging catalogues from individual detectors. Filenames are of the form atlas_er2_21h40-039d05_ugriz_finalSourceCat_NNNNNN.fits where NNNNNN is just an integer framesetID that represents the combination of observations used in the merger.

Source List and Catalogue Columns

A full description of the CASU source list columns is given at http://apm49.ast.cam.ac.uk/surveys-projects/vst/technical/catalogue-generation

Full descriptions of the WFAU bandmerged catalogue columns is given in Appendix 1. There are 155 columns per source. Flags uClass, gClass,... refer to the most probable morphological classification in each band where -1=stellar, +1=non-stellar, 0=noise and -2=borderline stellar. For description of other flags see the above link and http://osa.roe.ac.uk/www/ATLASDR5/ATLASDR5 TABLE atlasSourceSchema.

Each WFAU file is the catalogue of a "tile". But the source merging works only at the CCD detector/extension level. So what is supplied are really pseudo tile based merged catalogues from concatenating the 32 individual catalogues. The merging/seaming therefore still suffers from the CCD edge effects when there are large offsets between the filters so similar caveats apply as discussed at http://osa.roe.ac.uk/ - Known Issues

Duplicates can be partly avoided using (priOrSec =0 or priOrSec =framesetid) or indeed using the PRI-MARY_SOURCE keyword. However, while the latter will remove duplicates it will also have the effect of occasionally causing sources to miss measurements in some bands.

Acknowledgements

Full details of the VST OmegaCam instrument and data can be found at:- https://www.eso.org/sci/facilities/paranal/instruments/omegacam/doc/VST-MAN-OCM-23100-3110-2 7 1.pdf

The VST ATLAS survey was undertaken using ESO Program ID 177.A-3011(A,B,C,D,E.F,G,H,I,J,K,L,M,N) (see also Shanks et al 2015, MNRAS, 451, 4238).

According to the Data Access Policy for ESO data held in the ESO Science Archive Facility, all users are required to acknowledge the source of the data with appropriate citation in their publications.

Since processed data downloaded from the ESO Archive are assigned Digital Object Identifiers (DOIs), the following statement must be included in all publications making use of them:

 Based on data obtained from the ESO Science Archive Facility with DOI: https://doi.eso.org/ 10.18727/archive/66

Publications making use of data which have been assigned an archive request number (of the form XXXXXX) must include the following statement in a footnote or in the acknowledgement:

 Based on data obtained from the ESO Science Archive Facility under request number <request_number>.

Science data products from the ESO archive may be distributed by third parties, and disseminated via other services, according to the terms of the Creative Commons Attribution 4.0 International license. Credit to the ESO provenance of the data must be acknowledged, and the file headers preserved.

Appendix 1: WFAU bandmerged catalogue column descriptions

Column: Name; FITS data type; Description

- 1: IAUNAME; 31A; IAU Name (not unique)
- 2: sourceID; K; UID (unique over entire VSA via programme ID prefix) of this merged detection as assigned by merge algorithm
- 3: cuEventID; J; UID of curation event giving rise to this record
- 4: frameSetID; K; UID of the set of frames that this merged source comes from
- 5: ra2000; D; Celestial Right Ascension
- 6: dec2000; D; Celestial Declination
- 7: l; D; Galactic longitude
- 8: b; D; Galactic latitude
- 9: lambda; D; SDSS system spherical co-ordinate 1
- 10: eta; D; SDSS system spherical co-ordinate 2
- 11: priOrSec; K; Seam code for a unique (=0) or duplicated (!=0) source (eg. flags overlap duplicates).
- 12: umgPnt; E; Point source colour U-G (using aperMag3)
- 13: umgPntErr; E; Error on point source colour U-G
- 14: gmrPnt; E; Point source colour G-R (using aperMag3)
- 15: gmrPntErr; E; Error on point source colour G-R
- 16: rmiPnt; E; Point source colour R-I (using aperMag3)
- 17: rmiPntErr; E; Error on point source colour R-I
- 18: imzPnt; E; Point source colour I-Z (using aperMag3)
- 19: imzPntErr; E; Error on point source colour I-Z
- 20: umgExt; E; Extended source colour U-G (using aperMagNoAperCorr3)
- 21: umgExtErr; E; Error on extended source colour U-G
- 22: gmrExt; E; Extended source colour G-R (using aperMagNoAperCorr3)
- 23: gmrExtErr; E; Error on extended source colour G-R
- 24: rmiExt; E; Extended source colour R-I (using aperMagNoAperCorr3)
- 25: rmiExtErr; E; Error on extended source colour R-I

```
26: imzExt; E; Extended source colour I-Z (using aperMagNoAperCorr3)
```

- 27: imzExtErr; E; Error on extended source colour I-Z
- 28: mergedClassStat; E; Merged N(0,1) stellarness-of-profile statistic
- 29: mergedClass; I; Class flag from available measurements (1|0|-1|-2|-3|-9=galaxy|noise|stellar|probableStar|probableGalaxy|saturated)
- 30: pStar; E; Probability that the source is a star
- 31: pGalaxy; E; Probability that the source is a galaxy
- 32: pNoise; E; Probability that the source is noise
- 33: pSaturated; E; Probability that the source is saturated
- 34: eBV; E; The galactic dust extinction value measured from the Schlegel, Finkbeiner & Davis (1998) maps. This uses the correction given in Bonifacio, Monai & Beers (2000). This correction reduces the ex-
- tinction value in regions of high extinction (E(B-V)>0.1)
- 35: aU; E; The galactic extinction correction in the U band for extragalactic objects
- 36: aG; E; The galactic extinction correction in the G band for extragalactic objects
- $37\text{:}\ aR;\ E;\ The\ galactic\ extinction\ correction\ in\ the\ R\ band\ for\ extragalactic\ objects$
- 38: aI; E; The galactic extinction correction in the I band for extragalactic objects
- 39: aZ; E; The galactic extinction correction in the Z band for extragalactic objects
- 40: uPetroMag; E; Extended source U mag (Petrosian)
- 41: uPetroMagErr; E; Error in extended source U mag (Petrosian)
- 42: uAperMag3; E; Default point source U aperture corrected mag (2.0 arcsec aperture diameter)
- 43: uAperMag3Err; E; Error in default point/extended source U mag (2.0 arcsec aperture diameter)
- 44: uAperMag4; E; Point source U aperture corrected mag (2.8 arcsec aperture diameter)
- 45: uAperMag4Err; E; Error in point/extended source U mag (2.8 arcsec aperture diameter)
- 46: uAperMag6; E; Point source U aperture corrected mag (5.7 arcsec aperture diameter)
- 47: uAperMag6Err; E; Error in point/extended source U mag (5.7 arcsec aperture diameter)
- 48: uAperMagNoAperCorr3; E; Default extended source U aperture mag (2.0 arcsec aperture diameter)
- 49: uAperMagNoAperCorr4; E; Extended source U aperture mag (2.8 arcsec aperture diameter)
- 50: uAperMagNoAperCorr6; E; Extended source U aperture mag (5.7 arcsec aperture diameter)
- 51: uHlCorSMjRadAs; E; Seeing corrected half-light, semi-major axis in U band
- 52: uGausig; E; RMS of axes of ellipse fit in U
- 53: uEll; E; 1-b/a, where a/b=semi-major/minor axes in U
- 54: uPA; E; ellipse fit celestial orientation in U
- 55: uErrBits; J; processing warning/error bitwise flags in U
- 56: uAverageConf; E; average confidence in 2 arcsec diameter default aperture (aper3) U
- 57: uClass; I; discrete image classification flag in U
- 58: uClassStat; E; N(0,1) stellarness-of-profile statistic in U
- 59: uppErrBits; J; additional WFAU post-processing error bits in U
- 60: uSeqNum; J; the running number of the U detection
- 61: uXi; E; Offset of U detection from master position (+east/-west)
- 62: uEta; E; Offset of U detection from master position (+north/-south)
- 63: gPetroMag; E; Extended source G mag (Petrosian)
- 64: gPetroMagErr; E; Error in extended source G mag (Petrosian)
- 65: gAperMag3; E; Default point source G aperture corrected mag (2.0 arcsec aperture diameter)
- 66: gAperMag3Err; E; Error in default point/extended source G mag (2.0 arcsec aperture diameter)
- 67: gAperMag4; E; Point source G aperture corrected mag (2.8 arcsec aperture diameter)
- 68: gAperMag4Err; E; Error in point/extended source G mag (2.8 arcsec aperture diameter)
- $69: gAperMag6; \ E; \ Point \ source \ G \ aperture \ corrected \ mag \ (5.7 \ arcsec \ aperture \ diameter)$
- 70: gAperMag6Err; E; Error in point/extended source G mag (5.7 arcsec aperture diameter)
- 71: gAperMagNoAperCorr3; E; Default extended source G aperture mag (2.0 arcsec aperture diameter)
- 72: gAperMagNoAperCorr4; E; Extended source G aperture mag (2.8 arcsec aperture diameter)
- 73: gAperMagNoAperCorr6; E; Extended source G aperture mag (5.7 arcsec aperture diameter)
- 74: gHlCorSMjRadAs; E; Seeing corrected half-light, semi-major axis in G band
- 75: gGausig; E; RMS of axes of ellipse fit in G
- 76: gEll; E; 1-b/a, where a/b=semi-major/minor axes in G
- 77: gPA; E; ellipse fit celestial orientation in G
- 78: gErrBits; J; processing warning/error bitwise flags in G
- 79: gAverageConf; E; average confidence in 2 arcsec diameter default aperture (aper3) G
- 80: gClass; I; discrete image classification flag in G
- 81: gClassStat; E; N(0,1) stellarness-of-profile statistic in G

```
82: gppErrBits; J; additional WFAU post-processing error bits in G
```

- 83: gSeqNum; J; the running number of the G detection
- 84: gXi; E; Offset of G detection from master position (+east/-west)
- 85: gEta; E; Offset of G detection from master position (+north/-south)
- 86: rPetroMag; E; Extended source R mag (Petrosian)
- 87: rPetroMagErr; E; Error in extended source R mag (Petrosian)
- 88: rAperMag3; E; Default point source R aperture corrected mag (2.0 arcsec aperture diameter)
- 89: rAperMag3Err; E; Error in default point/extended source R mag (2.0 arcsec aperture diameter)
- 90: rAperMag4; E; Point source R aperture corrected mag (2.8 arcsec aperture diameter)
- 91: rAperMag4Err; E; Error in point/extended source R mag (2.8 arcsec aperture diameter)
- 92: rAperMag6; E; Point source R aperture corrected mag (5.7 arcsec aperture diameter)
- 93: rAperMag6Err; E; Error in point/extended source R mag (5.7 arcsec aperture diameter)
- 94: rAperMagNoAperCorr3; E; Default extended source R aperture mag (2.0 arcsec aperture diameter)
- 95: rAperMagNoAperCorr4; E; Extended source R aperture mag (2.8 arcsec aperture diameter)
- 96: rAperMagNoAperCorr6; E; Extended source R aperture mag (5.7 arcsec aperture diameter)
- 97: rHlCorSMjRadAs; E; Seeing corrected half-light, semi-major axis in R band
- 98: rGausig; E; RMS of axes of ellipse fit in R
- 99: rEll; E; 1-b/a, where a/b=semi-major/minor axes in R
- 100: rPA; E; ellipse fit celestial orientation in R
- 101: rErrBits; J; processing warning/error bitwise flags in R
- 102: rAverageConf; E; average confidence in 2 arcsec diameter default aperture (aper3) R
- 103: rClass; I; discrete image classification flag in R
- 104: rClassStat; E; N(0,1) stellarness-of-profile statistic in R
- 105: rppErrBits; J; additional WFAU post-processing error bits in R
- 106: rSeqNum; J; the running number of the R detection
- 107: rXi; E; Offset of R detection from master position (+east/-west)
- 108: rEta; E; Offset of R detection from master position (+north/-south)
- 109: iPetroMag; E; Extended source I mag (Petrosian)
- 110: iPetroMagErr; E; Error in extended source I mag (Petrosian)
- 111: iAperMag3; E; Default point source I aperture corrected mag (2.0 arcsec aperture diameter)
- 112: iAperMag3Err; E; Error in default point/extended source I mag (2.0 arcsec aperture diameter)
- 113: iAperMag4; E; Point source I aperture corrected mag (2.8 arcsec aperture diameter)
- 114: iAperMag4Err; E; Error in point/extended source I mag (2.8 arcsec aperture diameter)
- 115: iAperMag6; E; Point source I aperture corrected mag (5.7 arcsec aperture diameter)
- 116: iAperMag6Err; E; Error in point/extended source I mag (5.7 arcsec aperture diameter)
- 117: iAperMagNoAperCorr3; E; Default extended source I aperture mag (2.0 arcsec aperture diameter)
- 118: iAperMagNoAperCorr4; E; Extended source I aperture mag (2.8 arcsec aperture diameter)
- 119: iAperMagNoAperCorr6; E; Extended source I aperture mag (5.7 arcsec aperture diameter)
- 120: iHlCorSMjRadAs; E; Seeing corrected half-light, semi-major axis in I band
- 121: iGausig; E; RMS of axes of ellipse fit in I
- 122: iEll; E; 1-b/a, where a/b=semi-major/minor axes in I
- 123: iPA; E; ellipse fit celestial orientation in I
- 124: iErrBits; J; processing warning/error bitwise flags in I
- 125: iAverageConf; E; average confidence in 2 arcsec diameter default aperture (aper3) I
- 126: iClass; I; discrete image classification flag in I
- 127: iClassStat; E; N(0,1) stellarness-of-profile statistic in I
- 128: ippErrBits; J; additional WFAU post-processing error bits in I
- 129: iSeqNum; J; the running number of the I detection
- 130: iXi; E; Offset of I detection from master position (+east/-west)
- 131: iEta; E; Offset of I detection from master position (+north/-south)
- 132: zPetroMag; E; Extended source Z mag (Petrosian)
- 133: zPetroMagErr; E; Error in extended source Z mag (Petrosian)
- 134: zAperMag3; E; Default point source Z aperture corrected mag (2.0 arcsec aperture diameter)
- 135: zAperMag3Err; E; Error in default point/extended source Z mag (2.0 arcsec aperture diameter)
- 136: zAperMag4; E; Point source Z aperture corrected mag (2.8 arcsec aperture diameter)
- $137: z Aper Mag 4 Err; \ E; \ Error \ in \ point/extended \ source \ Z \ mag \ (2.8 \ arcsec \ aperture \ diameter)$
- 138: zAperMag6; E; Point source Z aperture corrected mag (5.7 arcsec aperture diameter)
- 139: zAperMag6Err; E; Error in point/extended source Z mag (5.7 arcsec aperture diameter)
- 140: zAperMagNoAperCorr3; E; Default extended source Z aperture mag (2.0 arcsec aperture diameter)

- 141: zAperMagNoAperCorr4; E; Extended source Z aperture mag (2.8 arcsec aperture diameter)
- 142: zAperMagNoAperCorr6; E; Extended source Z aperture mag (5.7 arcsec aperture diameter)
- 143: zHlCorSMjRadAs; E; Seeing corrected half-light, semi-major axis in Z band
- 144: zGausig; E; RMS of axes of ellipse fit in Z
- 145: zEll; E; 1-b/a, where a/b=semi-major/minor axes in Z
- 146: zPA; E; ellipse fit celestial orientation in Z
- 147: zErrBits; J; processing warning/error bitwise flags in Z
- 148: zAverageConf; E; average confidence in 2 arcsec diameter default aperture (aper3) Z
- 149: zClass; I; discrete image classification flag in Z
- 150: zClassStat; E; N(0,1) stellarness-of-profile statistic in Z
- 151: zppErrBits; J; additional WFAU post-processing error bits in Z
- 152: zSeqNum; J; the running number of the Z detection
- 153: zXi; E; Offset of Z detection from master position (+east/-west)
- 154: zEta; E; Offset of Z detection from master position (+north/-south)
- 155: PRIMARY_SOURCE; B; Primary source 1; secondary source 0