



Early Galaxy Formation and Evolution Through the Eyes of New Facilities




Brant Robertson
University of Arizona

Brief History of the Universe



Adapted from Robertson et al. *Nature*, **468**, 49 (2010).


Observational Facilities Over the Next Decade



Observations with ALMA, JWST, LSST, GMT/TMT/E-ELT, and WFIRST will drive astronomical discoveries over the next decade.

JWST


*hi res, NIR,
spectroscopy
space-based*



- 6.4m mirror in 18 segments
- 0.6-5 μ m NIRCам (30AB)
- 5-28.5 μ m MIRI
- R~100, 1000, 2700 @ 1-5 μ m

LSST

*wide area, optical,
ground-based*



- 8.4m with embedded tertiary
- 3.2 Gpixel camera; 60PB raw
- 18,000 deg² *ugrizy* AB=26.7
- 4x 9.6 deg² *ugrizy* AB>28

ALMA

*hi res, (sub)mm,
spectroscopy
ground-based*



- 66 x 12m antennae
- 0.3-9.6mm observations
- 15-18km baselines
- up to 0.005 arcsec resolution



WFIRST

*wide area, NIR,
R~600, space-based*

- 2.4m AFTA design from SDO
- 0.28 deg² ZYJHF detector
- 2,000 deg² YJHF AB=26.7
- 5-27.44 deg² 2b AB=27-29

TMT


*hi res, NIR,
spectroscopy,
ground-based*



- 30m segmented primary
- 15 arcmin FOV, 0.3-28 μ m
- NIR diffraction limited
- AO fed spectroscopy

UDF(12)

Beckwith et al. 2006
Koekemoer et al. 2013
Ellis et al. 2013
Illingworth et al. 2013



Multi-Cycle Treasury Programs




CANDELS: Cosmic Assembly Near-infrared Deep Extragalactic Legacy Survey

Grogin et al., ApJS, 197, 35 (2011)


Koekemoer et al., ApJS, 197, 36 (2011)

CLASH: The Cluster Lensing and Supernova Survey with Hubble
Postman et al., ApJS, 199, 25 (2012)



$z \sim 9$ Candidate; Zheng et al.,
Nature, 489, 406 (2012)






FRONTIER


FIELDS

- 6 Lensing clusters with ACS F435W, F606W, F814W and WFC3 F105W, F125W, F140W, and F160W (~28.8AB)
- Total of 840 orbits fully implemented
- Community funded to produce publicly-released gravitational lensing magnification maps.
- Observations complete for two clusters (Abell 2744, MACS J0416-2403)




Hubble Frontier Field Abell 2744
Hubble Space Telescope • ACS • WFC3

Continuing the High-z Galaxy Search with the Frontier Fields





Atek et al. arXiv:1311.7670




Zheng et al. arXiv:1402.6743

- DDT Hubble program to observe deeply gravitational lensing galaxy clusters.
- 70 ACS and 70 WFC3 orbits per cluster, sufficient for finding high-z galaxies.
- 16-18 $z > 6.5$ galaxies claimed in Abell 2744, more to come.

UV Luminosity Function @ $z \sim 7$



Physical Ingredients of Reionization



Ionization Fraction Evolution

$$\dot{Q}_{\text{HII}} = \frac{\dot{n}_{\text{ion}}}{\langle n_{\text{H}} \rangle} - \frac{Q_{\text{HII}}}{t_{\text{rec}}}$$

Ionizing photon production rate (density) $\rightarrow \dot{n}_{\text{ion}}$
 Comoving H density $\rightarrow \langle n_{\text{H}} \rangle$
 Recombination time $\rightarrow t_{\text{rec}}$

Ionizing Photon Density

$$\dot{n}_{\text{ion}} = f_{\text{esc}} \xi_{\text{ion}} \rho_{\text{UV}}$$

Escape fraction $\rightarrow f_{\text{esc}}$
 Ionizing photons per UV luminosity $\rightarrow \xi_{\text{ion}}$
 Comoving UV luminosity (SFR) density $\rightarrow \rho_{\text{UV}}$

Recombination Time

$$t_{\text{rec}} = [C_{\text{HII}} \alpha_{\text{B}}(T) (1 + Y_{\text{p}}/4X_{\text{p}}) \langle n_{\text{H}} \rangle (1 + z)^3]^{-1}$$

Clumping factor $\rightarrow C_{\text{HII}}$
 Case B recombination coefficient $\rightarrow \alpha_{\text{B}}(T)$
 Helium fraction $\rightarrow Y_{\text{p}}/4X_{\text{p}}$

Clumping Factor

$$C_{\text{HII}} \equiv \frac{\langle n_{\text{HII}}^2 \rangle}{\langle n_{\text{HII}} \rangle^2}$$


~1-6 (Pawlik et al. 2009)

Star Formation History of the Universe

Star formation history from UV (blue) and IR (red) observations, adopted from Madau & Dickinson (2014).

We can use this star formation history of the universe to compute the reionization history of the IGM.

BER, et al., ApJL, accepted
arXiv:1502.02024





Reionization History Implied by High- z Star Formation

IGM Neutral Fraction, shown with constraints from a variety of independent probes.

Galaxies with $L_{UV} > 0.001 L^*$ are sufficient to reionize the Universe by $z \sim 6$ (red band). (red band), consistent with other IGM neutrality probes.

Forced match to WMAP Thomson optical depth constraints (orange) requires earlier reionization with same assumptions, not consistent with IGM neutrality constraints.





Thomson Optical Depth to Electron Scattering

Recent Planck results suggest the Thomson optical depth to the CMB is $\tau \sim 0.065$, substantially less than found by WMAP.


The Thomson optical depth provided by the high- z IGM ionized by star forming galaxies is now fully consistent with the CMB constraints (Robertson et al. 2015).

BER, et al., ApJL, accepted
arXiv:1502.02024


Implications for High-Redshift Galaxy Observations with James Webb Space Telescope

The Thomson scattering signal in the CMB suggests the presence of high-redshift galaxies.

We expect $\sim 0.4\text{-}0.5$ $z > 10$ galaxies per square arcminute, detectable with JWST in the first deep exposures — ***we should find 5-10 $z > 10$ galaxies in the first year of JWST operations.***



Observations of Distant Galaxies with James Webb Space Telescope



The amazing array of capable instruments on JWST will enable truly synergistic surveys of the high-redshift universe!




NIRCam



NIRISS



MIRI







NIRSpec


- Spectroscopic confirmation of high-redshift galaxies with NIRSpec, MIRI, NIRISS, NIRCAM
- Spectral signatures of star formation rates and metallicity for high-redshift galaxies
- Clustering of high-redshift galaxies
- Multi-tier depth/area designs for distant galaxy surveys

JWST NIRCcam

- Simultaneous imaging in “red” (LW) and “blue” (SW) filters in two detector modules.
- Sensitivity is $\sim 29\text{AB}$ in 10^4s @ $2\mu\text{m}$
- Pixel scale in SW and LW arms give Nyquist sampling of PSF @ $2\mu\text{m}$ & $4\mu\text{m}$



Nebular Emission in High-z Galaxies




The rest-frame UV photometric data from HST and rest-frame optical data from *Spitzer* enable a fit of stellar population synthesis models to determine stellar masses.

Galaxies in certain redshift windows will have their rest-frame optical photometry affected by **nebular line emission**.

Nebular continuum emission can also affect rest-frame UV-optical photometry.

See, e.g., Schaerer & de Barros (2009, 2010); Ono et al. (2010); Robertson et al. (2010); Shim et al. (2011); de Barros et al. (2012); Stark et al. (2013), Labbe et al. (2013); de Barros, Schaerer, & Stark (2014); Smit et al. (2014, 2015)

Modeling Nebular Emission




1) Line Emission

$$L(H\beta) = 4.78 \times 10^{-13} (1 - f_{\text{esc}}) N_{\text{Lyc}} [\text{ergs s}^{-1}]$$

Tie additional lines (He, C, O, N, Mg, Ne, S, Ar) to H β luminosity (following Anders & Fritze-v. Alvensleben 2003).

Modeling Nebular Emission



1) Line Emission

$$L(H\beta) = 4.78 \times 10^{-13} (1 - f_{\text{esc}}) N_{\text{Lyc}} [\text{ergs s}^{-1}]$$

Tie additional lines (He, C, O, N, Mg, Ne, S, Ar) to H β luminosity (following Anders & Fritze-v. Alvensleben 2003).

Modeling Nebular Emission

1) Line Emission

$$L(H\beta) = 4.78 \times 10^{-13} (1 - f_{\text{esc}}) N_{\text{Lyc}} [\text{ergs s}^{-1}]$$

Tie additional lines (He, C, O, N, Mg, Ne, S, Ar) to H β luminosity (following Anders & Fritze-v. Alvensleben 2003).

2) Continuum Emission


$$L_{\nu} = \frac{\gamma_{\nu}}{\alpha_{\text{B}}} (1 - f_{\text{esc}}) N_{\text{Lyc}}$$

Nebular continuum emission proportional to Lyman continuum photon production.

Continuum Emission Coefficient


$$\gamma_{\nu} = \gamma_{\nu}^{\text{HI}} + \gamma_{\nu}^{2\nu} + \gamma_{\nu}^{\text{HeI}} \frac{n(\text{He}^{+})}{n(\text{H}^{+})} + \gamma_{\nu}^{\text{HeII}} \frac{n(\text{He}^{++})}{n(\text{H}^{+})}$$

Nebular Emission Coefficients



$$\gamma_{\nu} = \gamma_{\nu}^{\text{HI}} + \gamma_{\nu}^{2\nu} + \gamma_{\nu}^{\text{HeI}} \frac{n(\text{He}^+)}{n(\text{H}^+)} + \gamma_{\nu}^{\text{HeII}} \frac{n(\text{He}^{++})}{n(\text{H}^+)}$$

Standard SED Modeling of High-z Galaxies



The rest-frame UV photometric data from HST and rest-frame optical data from *Spitzer* enable a fit of stellar population synthesis models.

Best-fit Bruzual & Charlot 2003 model for stacked data:


Age = 4×10^8 yrs

SFR = $2 M_{\odot} / \text{yr}$

$M_{\star} = 1.25 \times 10^9 M_{\odot}$

SED Modeling of High-z Galaxies

Including Nebular Emission




Nebular emission models can fit the photometry equally well, or better than stellar only models. Even seemingly ridiculous ones.

Best-fit Bruzual & Charlot 2003 model + neb. emission ($f_{\text{esc}}=0$) for stacked data:

Age = 1.45×10^6 yrs
SFR = $36 M_{\text{sun}} / \text{yr}$
 $M_{\star} = 5.25 \times 10^7 M_{\text{sun}}$


Constraints on the nebular emission are going to be necessary to constrain tightly stellar mass at high-z.

Connecting Line Diagnostics in High Redshift Galaxies Across Wavelengths




What can we learn by combining observations from these two facilities?

High- z Star-Forming Galaxies are Strong UV-Optical Line Emitters




HST + Spitzer + NBLya

Ouchi et al. (2013)




Himiko $Ly\alpha$ Emitter @ $z\sim 6.6$


Far IR Emission(?) from High-z Galaxies



HST + Spitzer + NBLya





ALMA Continuum Map




ALMA [CII] Velocity Channel Maps

Himiko Ly α Emitter @ $z \sim 6.6$









See also Venemans et al. (2012), Carilli et al. (2013), Wagg et al. (2012), Carniani et al. (2013), Williams et al. (2014), Riechers et al. (2014), Rawle et al. (2014), De Brueck et al. (2014), Ferkinhoff et al. (2014), Hyunh et al. (2014), Swinbank et al. (2012, 2014), Wang et al. (2013), Willott et al. (2015)

CII 158 μm vs. Star Formation Rate



Ota et al. (2014)



Maiolino et al. (2015)

Collisional Excitation of CII 158 μm

Goldsmith et al. (2012), ApJS, 203, 13

Two-level model of collisional excitation of C^+ with H , H_2 , e^- .

Over a wide range of C^+ columns, emission increases with optical depth even when the line is optically thick.


Emission increases with kinetic temperature, and decreases with T of background radiation field (e.g., CMB).

Emission increases with ratio C_{ul}/A_{ul} , the collisional de-excitation vs. spontaneous emission rates.

Zero-excitation optical depth $\sim \text{C}^+$ column density:



$$\tau_0 = 7.49 \times 10^{-18} N(\text{C}^+)/\delta\nu \text{ (km s}^{-1}\text{)}$$

<https://github.com/brantr/c-plus>




Photodissociation Region Diagnostics

Kaufman et al. (1999), ApJ, 527, 795



Photodissociation Region Diagnostics


Kaufman et al. (1999), ApJ, 527, 795



Photodissociation Region Diagnostics

Kaufman et al. (1999), ApJ, 527, 795


n = density
 G_0 = ISRF




Photodissociation Region Diagnostics

Kaufman et al. (1999), ApJ, 527, 795

n = density
 G_0 = ISRF



LOTS of Information in the Rest-Frame Submillimeter!



SUMMARY : Early Galaxy Formation and Evolution Through the Eyes of New Facilities



- In the last 5 years, a new population of high-redshift galaxies have been discovered out to $z \sim 10$, informing us about the role of galaxies in reionization.
- Current uncertainties about the properties of high-redshift galaxies include the production rate of ionizing photons and the fraction that escape to ionize the IGM.
- Strong line emission in the UV-optical is inferred for many high-redshift galaxies, will be confirmed with JWST.
- The combined power of ALMA and JWST will teach us a lot about the ISM conditions of high-redshift galaxies and the nature of their stellar populations.