




The low-redshift universe as seen by the
Galaxy And Mass Assembly
survey

Jochen Liske
UHH





Bernhard Schmidt



Otto Heckmann





HERA


FLASH

DORIS

PETRA





NASA, ESA and the Hubble Heritage
(STScI/AURA)-ESA/Hubble Collaboration;
Davide de Martin and Robert Gendler



Planck Collaboration



Structure formation

$t = 4 \times 10^5$ yr
 $\Delta\rho/\rho = 10^{-5}$





$t = 1.37 \times 10^{10}$ yr
 $\Delta\rho/\rho = 10^9$

Gravitational instability and hierarchical build-up




De Lucia & Blaizot (2007)


Structure formation




Structure formation





Samushia et al. (2013)



Gravitational instability and hierarchical build-up




Increased density region Average density region





De Lucia & Blaizot (2007)

Gravitational instability and hierarchical build-up




Increased density region Average density region







NASA, ESA and the Hubble Heritage
(STScI/AURA)-ESA/Hubble Collaboration;
Davide de Martin and Robert Gendler




Bimodality

Renzini & Peng (2015)




Schawinski et al. (2014)

Mass distribution of galaxies





ellipticals



disk galaxies



irregular



The complexity of galaxies

Galaxy constituents

- Dark matter
- Stars and star clusters
- Molecular, atomic and ionised gas
- Dust
- Central SMBH
- IGM

Physical processes

- Gravitational collapse and evolution
- Gas hydrodynamics
- Star formation
- Stellar evolution
- Feedback
- Interaction with the environment
- ...

Outstanding issues

- Star formation efficiency and the nature of feedback as a function of halo mass
- Fuelling and cessation of star formation
- Roles of galaxy interactions and mergers versus in-situ processes
- Relative prevalence of disks and spheroids
- Mass-size relations of disks and spheroids
- Downsizing
- Co-evolution of central SMBH and their host galaxies
- ...

Complementary approaches

Observations

Statistical
investigations of
large samples
(surveys)

Level of detail
↔
Statistical power, completeness

Detailed studies
of small samples



Theory

Analytical, semi-analytical, numerical



The Galaxy And Mass Assembly survey

www.gama-survey.org

What is GAMA?

- A comprehensive survey of low-redshift ($z < 0.5$) galaxies to study galaxy evolution and cosmology
- GAMA = spectroscopic survey + alliance of imaging surveys

Key features

- $r < 19.8$ mag
- Area = 286 deg^2
- $N_{\text{gal}} = 270,000$
- Spectroscopy: 2 mag deeper than SDSS, multi-pass
- Imaging: near-complete wavelength coverage, $\sim 2x$ better resolution



The team

101 team members, 53 collaborators






GAMA survey regions


5 survey regions:
• 3 equatorial
• 2 southern

Total area = 286 deg²






Photometry: 1 nm – 1 m





GAMA spectroscopy



- 210 nights (4 FTEs!) of multi-object fibre spectroscopy using AAT/2dF+AAOmega
- Area: 286 deg² split over 5 survey regions
- Main sample: ~270k galaxies to $r < 19.8$ mag
- $R = 1300$, $370 < \lambda < 880$ nm
- $\langle z \rangle = 0.27$







GAMA spectroscopy


- 210 nights (4 FTEs!) of multi-object fibre spectroscopy using AAT/2dF+AAOmega
- Area: 286 deg^2 split over 5 survey regions
- Main sample: $\sim 270\text{k}$ galaxies to $r < 19.8$ mag
- $R = 1300, 370 < \lambda < 880$
- $\langle z \rangle = 0.27$






- DEEP2
- zCOSMOS
- GAMA
- 2dFGRS
- SDSS DR9
- 6dFGS











How does GAMA fit in?







Selection function management




- Dependence of redshift success on fibre position on the plate.
- Several possible causes, including:
 - Systematic errors in astrometry, field rotation, correction for atmospheric refraction, ...
 - Radial variation of apparent fibre diameter, focal ratio degradation, non-telecentricity, ...






Selection function management

- GAMA is a multi-pass survey by design.
- Tiling strategy is important! It affects the homogeneity of the incompleteness as well as survey efficiency.
- In GAMA: next tile placed where it most decreases local incompleteness.






Selection function management




➤ Smooth large-scale distribution of incompleteness.






Selection function management

Maintaining high redshift completeness in dense regions is crucial for the identification of groups and mergers.






Selection function management






Selection function management





Redshift precision...


From ~ 2000 duplicate observations:





...and accuracy

From ~ 2000 duplicate observations:





Technical papers

Spectroscopy

- Baldry et al. (2010)
- Robotham et al. (2010)
- Hopkins et al. (2013)
- Driver et al. (2011)
- Baldry et al. (2014)
- Liske et al. (2015)

input catalogue
tiling strategy
data reduction
survey procedures, DR1
redshift measurement code
end of survey report, QC, DR2

Photometry

- Hill et al. (2011),
Driver et al. (2016)
- Kelvin et al. (2012),
Häußler et al. (2013)
- Smith et al. (2011)
- Cluver et al. (2014)
- Liske et al. (2015)
- Driver et al. (2016)
- Prescott et al. (2016)
- Pacaud et al. (2016)
- Fotopoulou et al. (2016)

u – K aperture matched photometry
u – K Sérsic photometry
matching with Herschel sources
WISE photometry
GALEX photometry
integrating all photometry, PDR
radio fluxes
X-ray fluxes



Other

- Baldry et al. (2012)
- Hopkins et al. (2013)
- Loveday et al. (2012)
- Taylor et al. (2011)
- Robotham et al. (2011)
- Alpaslan et al (2014)
- Moffett et al. (2016)

local flow corrections
spectral line measurements
k-corrections
stellar masses
group catalogue
large-scale structure catalogue
morphologies




The stellar mass function




Baldry et al. (2012)



The stellar mass function



Peng et al. (2010)




Baldry et al. (2012)



BUT....


- Dust correction
- Issue is completely dominated by how you define *red*.
- A simple, hard colour cut is too simple.
- Not much evidence of a red population below $\log M^* \sim 9$.

Taylor et al. (2015)






Is the red faint upturn real?






GSMF by morphological type






GSMF by morphological type







GSMF by morphological type






GSMF by local environment







GSMF by local environment and z



McNaught-Roberts et al. (2014)




Model constraints



McNaught-Roberts et al. (2014)





Does the GSMF depend on LSS?







Does the GSMF depend on LSS?



Eardley et al. (2015)




Does the GSMF depend on LSS? No.






The GAMA group catalogue



- FoF algorithm to identify galaxy groups.
- FoF parameters carefully calibrated on mocks.
- $\sim 24,000$ groups in equatorial survey regions.
- 2754 groups with $N_{\text{fof}} > 4$



The GAMA group catalogue




- FoF algorithm to identify galaxy groups.
- FoF parameters carefully calibrated on mocks.
- ~24,000 groups in equatorial survey regions.
- 2754 groups with $N_{\text{fof}} > 4$

GAMA + KiDS \rightarrow weak lensing


Analysis of an initial joint GAMA+KiDS dataset covering 100 deg²

- The DM density profiles of group halos are well described by NFW
- Average halo masses \rightarrow scaling relations
- Provides constraints on feedback models on group scales






GAMA + KiDS → weak lensing




Viola et al. (2015)






Galaxy-galaxy weak lensing






Stellar-halo mass ratio in dense environments



Van Uitert et al. (in prep)




The masses of infalling sub-halos



Sifon et al. (2015)




The masses of infalling sub-halos






Panchromatic photometry






Panchromatic photometry



Driver et al. (2016)




Panchromatic photometry






Panchromatic photometry






Recalibrating SFR indicators




SFR density evolution






The effect of galaxy interactions on the SFR



Davies et al. (2015)




The effect of galaxy interactions on the SFR






The effect of galaxy interactions on the SFR






The effect of galaxy interactions on the SFR






The effect of galaxy interactions on the SFR



Davies et al. (2015)




The stellar mass dependence of close pairs






Stellar mass dependence of major merger rate






The evolution of the M^* close pair fraction





The role of mergers in building up stellar mass





You, too, can use GAMA data

► DR2

GAMA Data Release 2

The second GAMA data release (DR2) provides AAT/AAOmega spectra, redshifts and a wealth of ancillary information for 72,225 objects from the first phase of the GAMA survey (2008 - 2010, usually referred to as GAMA I). The DR2 web pages describe the data included in this release, and provide access to an SQL database as well as to the actual data (spectra and catalogues).


If you are using GAMA DR2 data in a publication then please cite the [DR2 paper \(Liske et al. 2015\)](#) and [acknowledge GAMA](#).

What is released?

The GAMA I survey extends over three equatorial survey regions of 48 deg^2 each (called G09, G12 and G15) and down to magnitude limits of $r < 19.4$ mag in G09 and G15, and $r < 19.8$ mag in G12. In DR2 we are releasing data for all GAMA I main survey objects with $r < 19.0$ mag (G09 and G12) or $r < 19.4$ mag (G15). Note that for G15 we are essentially releasing all GAMA I data. The total number of objects included in DR2 is 72,225. Of these, 70,726 objects (98%) have secure redshifts.

Details of the object selection for DR2:

- The qualifier 'GAMA I' refers to the fact that the objects for DR2 were selected from the input catalogue for the first phase of the GAMA survey (= GAMA I), see [Baldry et al. \(2010\)](#) for a detailed description of the GAMA I input catalogue.
- The qualifier 'main survey' refers to the fact that some targets were selected in different ways and for different reasons than those of the main GAMA survey. These so-called 'filler' targets were only observed when a fibre could not be allocated to a main survey target. Filler targets are not included in DR2.
- The r-band magnitude is the Petrosian r-band magnitude from SDSS DR6, corrected for Galactic extinction. This is the GAMA I selection magnitude.
- The three GAMA I survey regions are each $12 \times 4 \text{ deg}^2$ in size, for a total survey area of 144 deg^2 .





You, too, can use GAMA data

Current:

- DR2
- Panchromatic DR

Soon:

- DR3

www.gama-survey.org

