

spectra. The $J = J-0$ rotational line of HCN was claimed to be detected for the first time in Comet Kohoutek (1973 XII) by Huebner, Snyder, and Buhl. However, this detection could not be confirmed by any other group and subsequent searches in other comets were unsuccessful. The line was unambiguously detected in Comet P/Halley by three observing groups using three different radio telescopes in 1985-86. (Bockelée-Morvan et al., Schloerb et al.; Winnberg et al.). The $J = 3-2$ line of HCN (the $J = 2-1$ line lies at a frequency with strong atmospheric oxygen absorption) was then detected earlier this year in Comet Austin by a French and an American group. The present observation is the first successful detection of a comet by SEST and it has been confirmed by two other telescopes. Let us hope that SEST can continue to contribute to cometary spectroscopy.

Change at the ESO Schmidt Telescope

After a period of nearly 20 years in charge of the ESO Schmidt telescope, and after the successful completion of the taking of plates for the ESO Southern Surveys, *Hans-Emil Schuster* will hand over the reins to *Bo Reipurth*, staff astronomer at La Silla. Dr. Reipurth will take up his new operational responsibilities as of January 1, 1991, so please direct all Schmidt-related questions, enquiries, etc. to him after this date.

(continued from page 2)

Interestingly, in terms of atmospheric stability La Silla was found to be better than previously thought, with a measured median "seeing" of 0.76 arcsec. Paranal is better with a mean of 0.66 arcseconds, but of even greater importance is the fact that the number of clear nights of exceptional quality (seeing better than 0.5 arcsecond) is about 2.4 times higher on Paranal (16% of all nights) than on La Silla (7%).

The atmospheric conditions on Paranal will allow the VLT to take full advantage of its unique imaging and spectroscopic capabilities so that fainter and more distant objects can be observed than with any other telescope in the world. Moreover, when the VLT is supported by "adaptive optics", it will produce images that are almost as sharp as if it were in space. In the "interferometric" mode, when the light from

Announcement of the 3rd ESO/ST-ECF Data Analysis Workshop
ESO, Karl-Schwarzschild-Str. 2
Garching, Germany
April 22-24, 1991

The aim of the Workshop is to provide a forum for discussions of astronomical software techniques and algorithms. It is held annually during the spring (April/May) and centres on a different astronomical area each time. Due to available space, participation will be limited to 80 people. At the last Workshop several people could not be accommodated and we therefore recommend that you send in the corresponding participation and accommodation forms well before the deadline.

The topic for the 1991 Data Analysis Workshop will be analysis of direct imaging data. The scientific section of the meeting will consist of three sessions each starting with a main talk followed by presentation of papers of 5-10 minutes duration. The last day is reserved for general user meetings for MIDAS and ST-ECF.

The tentative agenda is:

Analysis of Direct Imaging Data

April 22: 14.00-18.00: Digital Filters
April 23: 9.00-12.30: Image Restoration
14.00-17.00: Decomposition techniques
17.00-18.00: European FITS Committee
April 24: 09.00-12.00: MIDAS user's meeting
12.00-13.00: European FITS Committee
14.00-17.30: ST-ECF user's meeting

Contributions on algorithms and techniques, e.g. removal of cosmic ray events on CCD's, digital transformations, deconvolution, decomposition of images and fitting techniques are especially welcome. We encourage people to present their work in these areas even if it is only ideas. After each introductory talk, we will have a more informal discussion where such contributions can be made. We also plan to have a poster session where people can present short contributions. Proceedings of the scientific sessions will be published.

The scientific organizing committee includes:

P. Grosbol (Chairman)	P. Benvenuti
L.B. Lucy	S. D'Odorico
D. Baade	R.H. Warmels

Contact address: Secretary of
Image Processing Group
European Southern Observatory
Karl-Schwarzschild-Str. 2
D-8046 Garching, Germany
EARN: DAW@DGAESO51
SPAN: ESO::DAW

the four 8.2-m telescopes is combined coherently (in the same phase), the resolving power of the VLT is further increased, so that even finer details can be seen. Under optimal circumstances, it should be possible to achieve a resolution of 0.0005 arcseconds. This would correspond to imaging 1 metre objects on the surface of the Moon.

Because of the extremely low atmospheric water vapour content in the Paranal region, probably the driest area on the surface of the Earth, this site is also highly suited for astronomical observations in the infrared and submillimetre wavelength regions.

The decision to place the VLT Observatory at Paranal implies that some years from now ESO will operate two, geographically separate observatories in Chile. In order to ensure the optimal functioning of both units, it will be necessary to adjust ESO's set-up in Chile.

The efficient running of the La Silla Observatory, on which so many European astronomers are dependent, will of course continue to have high priority, but it is expected that a certain streamlining will have to be made of the operations there.

The next step in the VLT programme will be to decide about the exact configuration of the four 8.2-metre telescopes and their enclosures. Several major contracts will be signed with European industry during the coming year, for instance for the construction of the mechanical structure of the giant telescopes and also the buildings which will be erected on Paranal.

The Editor

In its session on December 4, 1990, Council elected Professor Franco Pacini (Florence) as new President and Mr. Henrik Grage (Copenhagen) as Vice-President.