

# Formation and evolution of S0 galaxies: a SAURON case study of NGC 7332

Jesús Falcón-Barroso,<sup>1,2\*</sup> Reynier F. Peletier,<sup>1,3</sup> Eric Emsellem,<sup>3</sup> Harald Kuntschner,<sup>4</sup> Kambiz Fathi,<sup>1,3</sup> Martin Bureau,<sup>5</sup> Roland Bacon,<sup>3</sup> Michele Cappellari,<sup>2</sup> Yannick Copin,<sup>6</sup> Roger L. Davies<sup>7</sup> and Tim de Zeeuw<sup>2</sup>

<sup>1</sup>*School of Physics & Astronomy, University of Nottingham, Nottingham NG7 2RD*

<sup>2</sup>*Sterrewacht Leiden, Niels Bohrweg 2, 2333 CA, Leiden, the Netherlands*

<sup>3</sup>*CRAL, Observatoire de Lyon, F-69561 St-Genis Laval cedex, France*

<sup>4</sup>*Space Telescope European Coordinating Facility, European Southern Observatory, Karl-Schwarzschild-Str. 2, 85748 Garching, Germany*

<sup>5</sup>*Columbia Astrophysics Laboratory, 550 West 120th Street, 1027 Pupin Hall, MC 5247, New York, NY 10027, USA*

<sup>6</sup>*Institut de physique nucléaire de Lyon, 4 rue Enrico Fermi, 69622 Villeurbanne Cedex, France*

<sup>7</sup>*University of Oxford, Astrophysics, Keble Road, Oxford OX1 3RH*

Accepted 2004 February 2. Received 2004 January 26; in original form 2003 September 24

## ABSTRACT

We present SAURON integral-field observations of the S0 galaxy NGC 7332. Existing broad-band ground-based and *Hubble Space Telescope* (*HST*) photometry reveals a double-disc structure and a boxy bulge interpreted as a bar viewed close to edge-on. The SAURON two-dimensional stellar kinematic maps confirm the existence of the bar and inner disc but also uncover the presence of a cold counter-rotating stellar component within the central 250 pc. The  $H\beta$  and [O III] emission line maps show that the ionized gas has a complex morphology and kinematics, including both a component counter-rotating with respect to the stars and a fainter corotating one. Analysis of the absorption line-strength maps show that NGC 7332 is young everywhere. The presence of a large-scale bar can explain most of those properties, but the fact that we see a significant amount of unsettled gas, together with a few peculiar features in the maps, suggests that NGC 7332 is still evolving. Interactions as well as bar-driven processes must thus have played an important role in the formation and evolution of NGC 7332, and presumably of S0 galaxies in general.

**Key words:** galaxies: elliptical and lenticular, cD – galaxies: evolution – galaxies: formation – galaxies: individual: NGC 7332 – galaxies: kinematics and dynamics – galaxies: stellar content.

## 1 INTRODUCTION

The nature of S0 galaxies has been a matter of controversy since their appearance in the first morphological classification schemes (Hubble 1936; de Vaucouleurs 1959; van den Bergh 1960a, b) and many authors refer to them as normal spirals stripped of their gas (e.g. Sandage & Visvanathan 1978; Dressler 1980). Studies of S0 galaxies in clusters suggest that interactions between galaxies can lead to the loss of interstellar matter in the disc and convert spirals into S0s. Other theories favour gas loss through ram-pressure stripping from the intergalactic medium (Gunn & Gott 1972) or via galactic winds (Faber & Gallagher 1976), or assert that S0 galaxies are primordial galaxies which have entirely consumed their gas due to a high star-formation rate (Larson, Tinsley & Caldwell 1980).

When studying the formation of S0 galaxies, one is often drawn to examine the order of events during galaxy assembly. Early formation models favour the formation of the bulge before the disc, which is supported by several photometric studies (e.g. Caldwell 1983; Terndrup et al. 1994; Peletier & Balcells 1996; Peletier et al. 1999) and by the fact that S0s behave like ellipticals in the widely known scaling relations (i.e. colour–magnitude,  $Mg_2$ – $\sigma$  or fundamental plane). Numerical simulations raised the possibility of an alternative scenario (e.g. Pfenniger 1993) in which bulges are formed via secular evolution processes after the disc (e.g. Sellwood & Wilkinson 1993; Courteau, de Jong & Broeils 1996; Bureau & Freeman 1999; Carollo 1999; Merrifield & Kuijken 1999). Although such studies naturally focus on the two major components, i.e. the large-scale disc and the bulge, galaxies often show more structures, e.g. central discs, stellar kinematically decoupled components (KDCs) and multiple gaseous components. These are commonly observed in galaxies all along the Hubble sequence and are often used to emphasize the

\*E-mail: jfalcon@strw.leidenuniv.nl

**Table 1.** Properties of NGC 7332.

Parameter	Value	Source
Morphological type	S0 pec	de Vaucouleurs et al. (1991)
$M_R$ [mag]	-21.86	Balcells & Peletier (1994)
$B - R$ [mag]	1.40	Balcells & Peletier (1994)
Outer ellipticity	0.75	Andredakis et al. (1995)
Distance modulus [mag]	31.81	Tonry et al. (2001)
Distance scale [pc/arcsec]	111.6	

role of mergers during the formation of early-type galaxies (Toomre 1977; White & Rees 1978; Schweizer 1986; Franx & Illingworth 1988).

In the last 15 years, evidence for the presence of KDCs in early-type galaxies (e.g. NGC 4406, Bender 1988; NGC 4365, Wagner, Bender & Moellenhoff 1988) has usually been revealed through long-slit measurements. This can introduce a priori assumptions about the geometry of the central structures, and several works have tried to overcome this limitation by mapping the galaxies with multiple slits at different positions angles (Fisher, Illingworth & Franx 1994; Fisher 1997; Statler & Smecker-Hane 1999; Cappellari et al. 2002). However, the arrival of a new generation of integral-field units (IFUs) providing contiguous fields of view (FOV), such as OASIS (e.g. NGC 4621, Wernli, Emsellem & Copin 2002) or SAURON (Bacon et al. 2001; de Zeeuw et al. 2002), offers a much more efficient way to detect KDCs.

Early SAURON results (Bacon et al. 2001; de Zeeuw et al. 2002) reveal a variety of structures much richer than usually recognized in early-type galaxies. It is therefore important to use the information contained in these features to elucidate the key processes at work during galaxy formation. The galaxies concerned can become the ideal benchmarks against which to test the predictions of galaxy formation and evolution theories. NGC 7332, the galaxy studied in this paper, may be such a keystone.

NGC 7332, a boxy edge-on S0 galaxy,<sup>1</sup> was studied extensively in the past. The galaxy is mainly known for a bright counter-rotating and a faint corotating [O III] gas component with respect to the stars (Bertola, Buson & Zeilinger 1992; Fisher, Illingworth & Franx 1994). Those gas structures were confirmed by Plana & Boulesteix (1996), who mapped the H $\alpha$  emission via Fabry–Perot observations. NGC 7332 exhibits a rather regular broad-band morphology (Peletier & Balcells 1997) and its colours are somewhat bluer than those of elliptical galaxies of the same luminosity. Spectral analysis of the central regions reveals a luminosity-weighted age of about 6 Gyr (Gregg 1989; Vazdekis & Arimoto 1999).

In this paper, we use SAURON observations to characterize new and known features and improve our understanding of the morphology and dynamics of NGC 7332. We focus the analysis on the distribution and kinematics of the stellar and gaseous components, but we also discuss the corresponding SAURON line-strength maps. We present strong evidence for the presence of a central KDC and complex two-dimensional (2D) gas kinematics.

The main characteristics of NGC 7332 are provided in Table 1. Section 2 summarizes the observations discussed in this paper and their reduction. A photometric analysis and comparison with the literature is presented in Section 3. In Sections 4 and 5, we present and discuss the kinematic maps of stars and gas, and describe the

methods used to disentangle their respective spectral contributions. The line-strength analysis is presented in Section 6. We finally discuss possible formation and evolution scenarios for NGC 7332 in Section 7 and summarize our conclusions in Section 8.

## 2 OBSERVATIONS AND DATA REDUCTION

### 2.1 SAURON observations and data reduction

We observed NGC 7332 with the integral field spectrograph SAURON attached to the 4.2-m William Herschel Telescope (WHT) of the Observatorio del Roque de los Muchachos at La Palma, Spain, on 1999 October 13. We obtained four largely overlapping exposures of 1800 s each, producing more than 1500 spectra per exposure, including 146 sky spectra 1.9 arcmin away from the main field. SAURON delivers a spectral resolution of 4.3 Å full width at half maximum (FWHM) and covers the narrow spectral range 4810–5350 Å (1.1 Å pixel<sup>-1</sup>). The spatial sampling of individual exposures is performed by an array of square lenses of 0.94 arcsec, providing a FOV of  $\approx 33 \times 41$  arcsec<sup>2</sup>. The seeing at the time of the observations was stable at about 1.1 arcsec (FWHM). Flux, velocity and line-strength standard stars were observed during the same observing run for calibration purposes. Arc-lamp exposures were taken before and after each target frame for wavelength calibration. A tungsten lamp exposure was also taken at the end of the night in order to build the mask allowing us to extract the data from the CCD frames.


We followed the procedures described in Bacon et al. (2001) for the extraction, reduction and calibration of the data, using the specifically designed XSAURON software. The sky level was measured with the help of the dedicated sky lenses and subtracted from the target spectra. We merged the four individual extracted data cubes by spatially resampling the spectra on to a common squared grid. The dithering of individual exposures (with an original spatial sampling of  $0.94 \times 0.94$  arcsec<sup>2</sup>) enabled us to sample the merged data cube on to  $0.8 \times 0.8$  arcsec<sup>2</sup> pixels. The resulting merged mosaic FOV is  $\approx 34 \times 46$  arcsec<sup>2</sup> with a total of 1881 spectra. We then binned our final data cube using the Voronoi 2D binning algorithm of Cappellari & Copin (2003) to create compact bins with a minimum signal-to-noise ratio S/N of 60 per pixel. Most spectra have S/N in excess of 60, however, so that most of the original spatial elements remain unbinned (e.g.  $[S/N]_{\max} \approx 300$  in the central lens).

### 2.2 STIS spectroscopy and data reduction

We also retrieved Space Telescope Imaging Spectrograph (STIS) data from the *Hubble Space Telescope* (HST) archive at the Space Telescope Science Institute<sup>2</sup> (STScI) to obtain high spatial resolution spectroscopy of the centre of NGC 7332. The data form part of the observations of proposal ID 7566 by R. Green, covering the Ca II region between 8282 and 8835 Å. The G750M grating was used together with the  $52 \times 0.2$  arcsec<sup>2</sup> slit aperture, providing a dispersion of 0.56 Å pixel<sup>-1</sup> and a spatial sampling of 0.05 arcsec pixel<sup>-1</sup>. We retrieved the data reduced by the pipeline, making use of the best calibration files available at the time, but corrections for the effect of

<sup>1</sup> The galaxy is classified S0p in the RC3 (de Vaucouleurs et al. 1991) due to its boxy bulge (Sandage 1961).

<sup>2</sup> Based on observations made with the NASA/ESA *Hubble Space Telescope*, obtained from the data archive at the Space Telescope Science Institute. STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555.



**Figure 1.** SAURON FOV (thick box) superposed on the *I*-band image of Peletier & Balcells (1997) (left panel). The orientation is indicated in the bottom-left corner and is the same as for the SAURON maps shown in Figs 4–10. Contour levels are separated by 1 mag. The *I*-band major-axis surface brightness profile from Peletier & Balcells (1997) is shown in the right panel.

fringing and cosmic-ray removal were still required. Fringing corrections were applied using the prescriptions in Goudfrooij, Baum & Walsh (1997), whereas we used the `REDUCE` package (Cardiel 1999) to perform the cosmic ray rejection. A K0III star (HR7615) from the same data set was also retrieved and underwent the same reduction processes. It was used as a template for kinematical measurements.


### 2.3 *HST* and ground-based imaging

Imaging data were retrieved from several archives to perform a structural analysis. We retrieved a set of Wide-Field Planetary Camera (WFPC-1) images from the *HST* archive at STScI from the observing program of S. Faber (Prop. ID 2600, see Lauer et al. 1995). The observations were made using the F555W filter (*V*-band). The data set consists of three target frames with exposure times of 35, 140 and 140 s, respectively, and spatial sampling  $0.044 \text{ arcsec pixel}^{-1}$ . We combined the individual exposures and rejected cosmic rays with the routine `CRREJ`, available within the `IRAF`<sup>3</sup> package, thus resulting in a cosmic-ray-cleaned merged image. We also made use of a fully reduced ground-based Cousins *I*-band image from Peletier & Balcells (1997), taken with the Isaac Newton Telescope (INT) located in the Observatorio del Roque de los Muchachos in La Palma, Spain. The exposure time was 200 s and the seeing 1.0 arcsec. We overlay in Fig. 1 the contours from this *I*-band image and the FOV and contours of the SAURON reconstructed total intensity map (obtained by summing the data in wavelength). Finally, the photometric analysis performed by Fisher et al. (1994) from an *R*-band image, taken in 1.3-arcsec seeing, was also used.

## 3 PHOTOMETRIC ANALYSIS

We analysed the WFPC-1 data in order to study morphological substructures in NGC 7332. A comparison of our analysis with that of Fisher et al. (1994) from an *R*-band image is shown in Fig. 2 and the agreement in the overlapping region is excellent. This comparison is possible because the colour gradients in this galaxy are small (Peletier & Balcells 1997) and the emission lines can be neglected (e.g.  $H\alpha$  and  $[N \text{ II}]$  in the *R*-band image).

The isophotal analysis of the *HST* image reveals a number of maxima and minima in the  $c_4$  parameter, which describes deviations of the isophotes from pure ellipses (positive  $c_4$  indicates discy



**Figure 2.** Photometric analysis of NGC 7332. The ratio of the WFPC-1 image and a model image from our best ellipse fit is shown in the top-left panel. Position angle (PA), ellipticity ( $\epsilon$ ) and boxiness ( $c_4$ ) profiles along the major-axis radius ( $r$ ) are shown in the other panels. Solid circles represent our WFPC-1 analysis whereas open squares show the profiles obtained by Fisher et al. (1994). The formal errors on the different parameters (on the WFPC-1 data) are 0.60, 0.005 and 0.004 for the PA,  $\epsilon$  and  $c_4$ , respectively.


isophotes and negative  $c_4$  boxy isophotes; see e.g. Carter 1978). Positive values are present in the inner 0.5 arcsec but may be strongly influenced by the halo of the pre-COSTAR WFPC-1 point spread function (PSF). Positive  $c_4$  values are also observed in the region  $1.5 \lesssim r \lesssim 7$  arcsec, where the maximum reaches about 3 per cent. The isophotes are then elliptical or slightly boxy at intermediate radii ( $7 \lesssim r \lesssim 12$  arcsec) and again strongly discy at large radii ( $r \gtrsim 12$  arcsec). A complementary representation is shown in Fig. 2 (top-left panel), where we divide our image by a 2D model from the best-fitting pure elliptical isophotes. This confirms the  $c_4$  behaviour and highlights the excess of light above the fitted elliptical model in the central 1–7 arcsec of the galaxy. We associate this with a central disc whose major axis coincides with that of the galaxy, at a position angle (PA) of  $160^\circ$ . The presence of such a central disc was already reported by Seifert & Scorza (1996). There is also evidence for a secondary disc at larger radii ( $\gtrsim 12$  arcsec), revealed by an excess of light and a positive  $c_4$  parameter (Fig. 2).

Another noteworthy feature is the position angle twist observed from the centre to the outer regions of the galaxy. The magnitude of the twist is about  $5^\circ$ . This PA change is most likely a projection effect due to a point symmetric structure (i.e. spiral-like structure) showing up in the outer disc, viewed at a large but not perfectly edge-on inclination.

## 4 STELLAR KINEMATICS

We measured the SAURON stellar kinematics of NGC 7332 by fitting its spectra with a linear combination of single-age, single-metallicity population models from Vazdekis (1999) convolved with a Gauss–Hermite expansion. The best-fitting parameters are determined by chi-squared minimization in pixel space using the penalized pixel fitting (pPXF) method by Cappellari & Emsellem (2004), which is implemented as part of the XSAURON software developed at the Centre de Recherche Astronomique de Lyon (CRAL). We computed the uncertainties by means of Monte Carlo simulations

<sup>3</sup> IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.



**Figure 3.** Comparison of kinematic profiles of NGC 7332 from SAURON (solid symbols) and Fisher et al. (1994, open symbols) along two position angles (PA = 155° and 65°).

(100 realizations) and estimate them to be  $5 \text{ km s}^{-1}$  on average for the mean velocity,  $11 \text{ km s}^{-1}$  for the velocity dispersion and 0.03 for  $h_3$  and  $h_4$  for  $S/N = 60$  per pixel. This procedure traces only errors due to sampling, spectral range and noise, but not those introduced by a potential template mismatch. We minimize template mismatch, however, by using an optimal template (see Section 4.1 below). The stellar kinematics were also measured in the inner arcseconds using the STIS data and the code of van der Marel & Franx (1993), restricting the fit to a Gaussian without higher-order terms because of the poor data quality.

#### 4.1 Measurement of emission-free stellar kinematics

NGC 7332 presents the following emission lines in the rest wavelength range covered by SAURON (4790–5350 Å):  $H\beta$  (4861 Å),  $[O \text{ III}]$  (4959 and 5007 Å) and the  $[N \text{ I}]$  doublet (5197.9 Å and 5200.4 Å). Caution is thus required when deriving the stellar kinematics. To obtain a clean stellar spectrum, we followed the detailed prescriptions described below.

We first build a library of synthetic stellar spectra with different ages and metallicities from the library of Vazdekis (1999), convolved to the SAURON instrumental resolution (rendered uniform over the

FOV). For each target spectrum, we then perform the following steps:

(i) We measure the stellar kinematics using pPXF in the wavelength range 4830–5280 Å, avoiding the regions with nebular emission (e.g.  $H\beta$ ,  $[O \text{ III}]$  and  $[N \text{ I}]$  lines). This procedure provides the best template fit to the full wavelength range.

(ii) The subtraction of the best template fit from the original data set results in a ‘pure emission line’ spectrum that is used to extract the gas kinematics. For this, we fit the emission spectrum approximating each line with a Gaussian. The Gaussian fit is then removed from the original data, yielding an ‘emission-line-free’ spectrum from which clean line-strength indices can be derived.


An illustration of this procedure for NGC 7332 is presented in Section 5.

In order to check the accuracy of our results, we compared them with the best available long-slit stellar kinematics of NGC 7332 (Fisher et al. 1994). In Fig. 3, the velocity and velocity-dispersion profiles along two perpendicular position angles, PA = 155° and PA = 65°, are shown. They are consistent within the errors.


#### 4.2 A kinematically decoupled component and central disc in NGC 7332

As shown in Fig. 4, the stellar kinematics of NGC 7332 display a rather smooth velocity field with rotation along the major axis, but it also shows some peculiarities. One is a KDC reported here for the first time ( $r \lesssim 2.5 \text{ arcsec}$ ) and revealed by an ‘S’-shaped zero-velocity contour in the SAURON stellar velocity field, emphasized in the enlarged version provided in Fig. 5 (bottom-left panel). We also present a comparison of the STIS spectroscopy and SAURON data along PA  $\approx 160^\circ$  (the apparent major axis in the inner parts; see Fig. 6). At the spatial resolution of SAURON, we only detect a weak central counter-rotating structure, but the signature is clear in the STIS data, with an amplitude of about  $30 \text{ km s}^{-1}$  in the central 2 arcsec.

In order to emphasize these central features, we have modelled the SAURON velocity field on large scales with a simple exponential disc (Freeman 1970), excluding the central 2.5 arcsec from the fit to avoid being affected by the KDC. The result is shown in Fig. 5. The best model was forced to have PA = 160°, consistent with the photometric major axis as measured in Section 3 and from the SAURON reconstructed total intensity image. The residual




**Figure 4.** Stellar kinematics of NGC 7332 from SAURON. (a) Reconstructed total intensity ( $\text{mag arcsec}^{-2}$ , arbitrary zero point), (b) mean velocity ( $\text{km s}^{-1}$ ), (c) velocity dispersion ( $\text{km s}^{-1}$ ), (d) and (e) Gauss–Hermite moments  $h_3$  and  $h_4$ . The SAURON spectra have been spatially binned to a minimum  $S/N$  of 60 by means of the Voronoi 2D binning algorithm of Cappellari & Copin (2003). Isophotes from the reconstructed total intensity image are overlaid on all maps in  $1 \text{ mag arcsec}^{-2}$  steps.




**Figure 5.** Fit to the SAURON velocity field of an exponential disc model (Freeman 1970). Top panels: original velocity field as shown in Fig. 4 (left) and fitted model (right). Bottom panels: zoom on the SAURON velocity field with the zero isovelocity curve superimposed (solid line; left) and the same region after subtraction of the best-fitting model (residual  $V - V_{\text{model}}$ ; right). All the maps are in units of  $\text{km s}^{-1}$ . The area within the dashed circle was excluded from the fit.

image (Fig. 5, bottom-right panel) highlights the KDC component: its angular momentum projected on to the minor axis of NGC 7332 is negative, justifying its identification as a counter-rotating component (CRC). The kinematic major axis of this CRC seems, however, to be tilted by  $\approx 20^\circ$  with respect to that of the main galaxy.

We note that there is only very weak evidence for a flattening of the velocity profile in the central arcseconds along  $\text{PA} = 155^\circ$  (see Fig. 3;



**Figure 6.** Comparison of the SAURON (solid symbols) and STIS (open symbols) velocity profiles at a position angle of  $160^\circ$  (the major axis). The SAURON error bars are, in some cases, smaller than the symbol size.




**Figure 7.** Stellar kinematic major-axis profiles of NGC 7332 from SAURON ( $\text{PA} = 160^\circ$ ). The radial velocity and velocity dispersion are  $\text{km s}^{-1}$ .

Section 4.1) and we cannot find any conclusive evidence for the presence of counter-rotation at this PA. It is, however, clearly seen at  $\text{PA} = 160^\circ$  (the major axis; see Fig. 6); and although the detection of this structure is strongly dependent on the spatial resolution, it again illustrates the advantage of IFUs over long-slit spectrographs.

The velocity-dispersion map shown in Fig. 4 reveals a central dip of about  $10 \text{ km s}^{-1}$  along the major axis of NGC 7332 (see also Fig. 7), in contrast with the traditional peak found at the centre of the bulges of most early-type galaxies. This drop in velocity dispersion is on the same scale as the KDC ( $r \lesssim 2.5 \text{ arcsec}$ ; see Fig. 4) and may well be associated with it.

In addition to the photometric evidence presented in Section 3, there are also kinematic features supporting the presence of a central disc on a larger scale than the KDC. In particular, the Gauss-Hermite moment  $h_3$  ( $\approx 0.1$  in amplitude) is strongly anticorrelated with  $V$  in the region where the central disc is postulated. The superposition of a fast-rotating disc to a moderately rotating spheroid produces a similar asymmetry in the line-of-sight velocity distributions (LOSVDs).

We have therefore discovered a remarkably unusual situation in the core of NGC 7332, with a stellar KDC ( $r \lesssim 2.5 \text{ arcsec}$ ) and a larger central disc ( $r \lesssim 7 \text{ arcsec}$ ).



**Figure 8.** Optimal template fitting for NGC 7332 in four different locations: the KDC (circle), main disc (upward triangle), minor axis (square) and bar (downward triangle). Red lines represent the best-fitting stellar templates (see Section 4.1) of the underlying galaxy spectra (black lines). Residual spectra are shown underneath. The central panel shows the [O III] intensity distribution (Fig. 9) with isophotes from the reconstructed total intensity image (Fig. 4) overlaid in  $1 \text{ mag arcsec}^{-2}$  steps.


## 5 GAS KINEMATICS

NGC 7332 is best known for its peculiar gas structure, namely the presence of two kinematically decoupled gas components, as observed by Bertola et al. (1992), Fisher et al. (1994) and Plana & Boulesteix (1996). Fisher et al. (1994) measured the [O III] emission using several slits along different position angles, whereas Plana & Boulesteix (1996) obtained complete 2D emission maps using the H $\alpha$  line (Fabry–Perot observations). Here, we are able to map three different emission lines within the SAURON wavelength range (H $\beta$ , [O III], [N I]).

We obtained a pure emission-line data cube using the procedure described in Section 4.1. An illustration of the method is shown in Fig. 8 for four different locations in the SAURON FOV. The fitted synthetic spectra (red lines) are overlaid on the SAURON spectra of NGC 7332 (black lines) and the differences are shown underneath. The residuals are small outside the spectral regions

where emission lines are expected to contribute, with oscillations just slightly above that expected from the corresponding noise level. As shown in Fig. 9, [O III] is strong everywhere in the galaxy. The H $\beta$  line, although present, is much fainter than [O III]. We measured the H $\beta$  and [O III] kinematics by fitting a single Gaussian to each of the three lines simultaneously, which relies on the assumption that all the lines have the same velocity and velocity dispersion. We have also assumed a 1:2.96 ratio for the [O III] to H $\beta$  ratio (Osterbrock 1989) but left the [O III] to H $\beta$  ratio free. In our analysis, these assumptions seem to hold everywhere in the field. Located on the wing of the Mg *b* absorption line (5170 Å), the [N I] doublet is very difficult to detect. Our analysis reveals negligible amounts of [N I] emission within the SAURON FOV so we have not corrected the original spectra for its presence.

In Fig. 9, we present the H $\beta$  and [O III] intensity maps as well as their kinematics in the form of mean radial-velocity and velocity-dispersion maps. The intensity distribution of the H $\beta$  and [O III]



**Figure 9.** H $\beta$  and [O III] gas distribution and kinematics from SAURON. The intensities are in  $\text{mag arcsec}^{-2}$  (arbitrary zero point) while the velocity and velocity dispersion are in  $\text{km s}^{-1}$ . Isophotes from the reconstructed total intensity image (Fig. 4) are overlaid on all maps in  $1 \text{ mag arcsec}^{-2}$  steps.

lines have similarities, although the  $H\beta$  intensity is much fainter. The detection of  $H\beta$  emission at the centre is somewhat marginal (compared to the bright underlying stellar contribution). We find a bright counter-rotating ionized gas component and weak traces of a much fainter corotating one. We were able to obtain [O III] intensity values for the main counter-rotating component easily but the second component is much more difficult to isolate, not only because of its low relative contribution but also because of the limited spectral resolution of SAURON (see Fig. 8). We could thus only trace it in a small region of our field, and decided not to attempt to derive its velocity field. There is, however, no doubt that this second component exists, as emphasized by Fisher et al. (1994) and Plana & Boulesteix (1996).

The emission-line maps we have obtained for NGC 7332 show a complex structure. The bulk of the [O III] emission is clearly misaligned with respect to the main stellar disc and shows strong departures from an axisymmetric distribution. We detect a long arc-like filament in the [O III] distribution on the north-west side of NGC 7332, connecting on to the major axis of the galaxy at the northern limit of the SAURON FOV. The velocity and velocity-dispersion maps of the gaseous component also show strong non-axisymmetric kinematics and we find an [O III] emitting region isolated in velocity ( $\approx 350 \text{ km s}^{-1}$  from systemic) about 7 arcsec south-west of the galaxy centre.

Masking out all marginal [O III] detections (essentially all flux below  $-3$  in Fig. 9) and the high-velocity material mentioned above, the [O III] map then exhibits a bar or spiral-like structure within the central 10 arcsec. This structure is similar to, but less contrasted than, the one found in the other nearly edge-on lenticular galaxy NGC 3377 (Bacon et al. 2001). Assuming that NGC 7332 is close to edge-on, at an inclination of  $\approx 75^\circ$  (Andredakis, Peletier & Balcells 1995), then the [O III] most likely has a substantial vertical extension and is simply seen in projection against the disc. If, on the other hand, NGC 7332 is more face-on, then this structure could indeed be in the equatorial plane.

## 6 STELLAR POPULATIONS

The many distinct kinematic components identified in the previous sections raise the question of the sequence of events leading to the current structure of NGC 7332. Valuable information can be obtained by investigating the stellar populations over the full FOV of SAURON. The measurement of absorption line-strengths together with predictions of stellar population models can be used to infer the luminosity-weighted age and metallicity of the stellar populations (e.g. González 1993; Worthey et al. 1994; Kuntschner & Davies 1998).

We used our flux-calibrated data cube to measure several line-strength indices available in the SAURON wavelength range (i.e.  $H\beta$ , Fe5015, Mg *b* and Fe5270). These indices were then calibrated on to the Lick/IDS system (Worthey et al. 1994) following the procedures outlined in Kuntschner (2000). Specifically, we observed 73 stars in common with the Lick/IDS system to establish the small systematic offsets for each index (see also Worthey & Ottaviani 1997). We also estimated the velocity-dispersion corrections, to account for the broadening of the spectral features by the LOSVDs, using a subset of model spectra by Vazdekis (1999) broadened to the Lick/IDS resolution.

The Lick/IDS Fe5270 index cannot be fully mapped with SAURON due to the varying bandpass over its FOV (de Zeeuw et al. 2002). In order to maximize the available FOV, a new index was defined (Fe5270<sub>s</sub>) which measures the same spectral feature, but has a reduced spectral coverage in the red pseudo-continuum band. This

new index was then converted on to the original Lick/IDS system via the empirical, linear relation

$$\text{Fe5270} = 1.26\text{Fe5270}_s + 0.12. \quad (1)$$

The  $1\sigma$  standard deviation of this empirical calibration is  $\pm 0.05 \text{ \AA}$  for the Fe5270 index. More details on the SAURON line-strength system will be given in a forthcoming paper of the main SAURON series that applies to the full SAURON survey.

We measured the indices before and after applying our emission-subtraction procedure (Section 4.1). As can be seen in the line-strength maps shown in Fig. 10, the corresponding corrections are large in the outer parts due to the substantial contamination by the emission lines (mostly  $H\beta$  and [O III]). The emission-corrected line-strength values do somewhat depend on the combination of single-age, single-metallicity population models used to build the optimal template and fit the raw data. But although this procedure is critical for the removal of the  $H\beta$  contamination from the  $H\beta$  index, it is negligible for the removal of the [O III] contamination from the Fe5015 index, and the Mg *b* and Fe5270 indices are totally unaffected. The maximum error in the  $H\beta$  index introduced by using different sets of optimal template libraries is  $0.08 \text{ \AA}$  ( $1\sigma$ ).


After the emission-subtraction process, we find that  $H\beta$  remains nearly constant (within the errors) over the whole SAURON FOV. Since the fitting procedure works on the spectra completely independently from one another, it is unlikely that a flat  $H\beta$  map could be obtained by an inaccurate subtraction of the emission. The Mg *b*, Fe5015 and Fe5270 maps show a steep increase in line-strength within 5 arcsec of the centre (see also Fig. 11), but outside of this inner region the metal line-strengths are approximately constant over the FOV. There is, however, some evidence that the values of the Mg *b*, Fe5015 and Fe5270 indices decrease slightly at  $r \approx 10$  arcsec and increase again at the outermost radii along the major axis (see Fig. 10 and Fig. 11).

The overall line-strength distribution is sufficiently uniform that none of the galaxy components identified in the previous sections (i.e. the KDC, inner disc, bar and outer discs) can be clearly identified in these maps. Specifically, the flat  $H\beta$  index suggests a rather uniform age across the entire FOV covered by SAURON. If the (luminosity-weighted) age of the bulge or the central disc in NGC 7332 were much different from that of the main body, it should be seen in the  $H\beta$  map ( $H\beta$  is sensitive to age and independent of metallicity changes to first order; see Worthey et al. 1994).


Index–index diagrams such as  $H\beta$  versus [MgFe],<sup>4</sup> together with stellar population model predictions, are often used to estimate luminosity-weighted ages and metallicities of galaxies. This combination of indices is almost insensitive to non-solar abundance ratios which can otherwise significantly affect age and metallicity estimates (e.g. Trager et al. 2000; Kuntschner et al. 2001; Thomas, Maraston & Bender 2003). Here, the use of the Fe5270 and Mg *b* indices as metallicity indicators is unfortunately limited by the restricted wavelength range of SAURON. In order to minimize the effects of non-solar abundance ratios, we have thus defined a new index, [MgFe52]  $\equiv (0.62\text{Mg}b + \text{Fe5270})/2.0$ , using the model predictions of Thomas et al. (2003).

In Fig. 12, we present the results of our analysis by averaging  $1.6 \times 1.6 \text{ arcsec}^2$  regions (i.e.  $2 \times 2$  lenslets) around the four key positions shown in Fig. 8. The estimated luminosity-weighted age for all regions is  $5 \pm 2 \text{ Gyr}$ . The metallicity of the central area is about 2 times solar while the outer areas

<sup>4</sup> [MgFe]  $\equiv \sqrt{\text{Mg}b \frac{(\text{Fe5270} + \text{Fe5335})}{2}}$




**Figure 10.** SAURON  $H\beta$ , Fe5015, Mg  $b$  and Fe5270 line-strength maps on the Lick/IDS system (Worthey et al. 1994). Top row: raw indices. Bottom row: emission-corrected indices. Isophotes from the reconstructed total intensity image (Fig. 4) are overlaid on all maps in 1 mag arcsec $^{-2}$  steps.



**Figure 11.** SAURON emission-corrected line-strength major-axis profiles ( $H\beta$ , Mg  $b$ , Fe5015 and Fe5270) on the Lick/IDS system (Worthey et al. 1994). The error bars represent Poisson errors.

show metallicities between 0.35 and 1 times solar. These results are consistent with others found in the literature. Using a similar line-strength analysis, the nucleus of NGC 7332 has been reported to host a population of luminosity-weighted age between 4 and 6 Gyr depending on the authors (Vazdekis & Arimoto 1999; Terlevich & Forbes 2002). The colour gradients are also



**Figure 12.**  $H\beta$  versus  $[MgFe52]'$  equivalent width diagram. Symbols represent measurements from the clean data cube at the four positions shown in Fig. 8: the KDC (circle), main disc (upward triangle), minor axis (square) and bar (downward triangle). The error bar in the bottom-left corner shows the  $1\sigma$  uncertainty introduced by the emission subtraction process. The error bars shown for the data points represent Poisson errors.

**Table 2.** SAURON parameters for NGC 7332.

Parameter	Value	Units
Stellar heliocentric systemic velocity	$1206 \pm 5$	$\text{km s}^{-1}$
Gas heliocentric systemic velocity	$1206^a \pm 7$	$\text{km s}^{-1}$
Stellar $V_{\text{max}}$	$123^b \pm 5$	$\text{km s}^{-1}$
Extent of the KDC	$\sim 2.5$	arcsec
Extent of the central disc	$\sim 7.0$	arcsec
Mass of counter-rotating ionized gas	$\sim 1.5$	$10^5 M_{\odot}$
EW( $H\beta$ )	$2.23^c \pm 0.09$	$\text{\AA}$
EW(Mg $b$ )	$3.45^c \pm 0.18$	$\text{\AA}$
EW([MgFe52])	$2.53^c \pm 0.07$	$\text{\AA}$

<sup>a</sup>From the  $H\beta$  and [O iii] lines fitted simultaneously.

<sup>b</sup>Within the SAURON field.

<sup>c</sup>Central aperture [ $1.6 \times 1.6 \text{ arcsec}^2$ ].

small in the disc and bulge (Peletier & Balcells 1997), consistent with our finding of an approximately constant age across the SAURON FOV.

The young luminosity-weighted age of  $5 \pm 2$  Gyr detected in our analysis clearly shows that NGC 7332 has experienced relatively recent star formation. Furthermore, the uniformity of the  $H\beta$  map suggests that this star formation affected all regions of the galaxy probed by our integral-field instrument.

## 7 DISCUSSION

In this paper, we studied a galaxy that has sometimes been labelled peculiar. Using integral-field spectroscopy, we were better able to constrain the stellar kinematics, the ionized gas distribution and kinematics, and the stellar populations of NGC 7332. In turn, these can now be used to constrain its evolutionary history better. A summary of some key numbers extracted from the data is given in Table 2.

### 7.1 The presence of a bar

A number of photometric features described in Section 3 are consistent with NGC 7332 being a barred galaxy viewed close to edge-on. Indeed, soon after they form, bars buckle and settle with an increased thickness, appearing boxy-shaped when seen end-on (i.e. along the bar major axis) and peanut-shaped when seen side-on (i.e. along the bar minor axis; see e.g. Combes et al. 1990 and Raha et al. 1991 for  $N$ -body simulations; Bureau & Freeman 1999 and Merrifield & Kuijken 1999 for observations). The boxiness of the isophotes does not show properly in the  $c_4$  profile of Fig. 2 because the isophotes are simultaneously boxy (at large galactic heights) and discy (along the equatorial plane), due to the superposition of the boxy bar and an inner disc. The boxiness is, however, obvious in the image shown in Fig. 1. The plateau in the major-axis surface brightness profile at intermediate radii found by (Seifert & Scorza 1996) and confirmed by our  $I$ -band image (Fig. 1, right panel) is also characteristic of bars viewed nearly edge-on (see e.g. Lütticke, Dettmar & Pohlen 2000; Bureau & Athanassoula 2004). NGC 7332 finally exhibits a double-disc structure, which can be interpreted as the result of secular evolution due to the presence of a bar (see, e.g. van den Bosch & Emsellem 1998 and references therein).

Although NGC 7332 is close to edge-on, it is still possible to use its (projected) kinematics to verify the existence of the large-scale thick bar suggested by the photometry (see Section 3; Bureau & Athanassoula 2004). The extracted major-axis kinematic pro-

files of NGC 7332 (Fig. 7; but see also Fisher et al. 1994) show kinematic features commonly seen in early-type barred galaxies: a ‘double-hump’ velocity profile and a central (local) minimum in the velocity dispersion. Double-hump velocity profiles are observed in dissipationless simulations of barred discs (Bureau & Athanassoula 2004) and in a number of galaxies with so-called ‘double-disc’ structures (e.g. Seifert & Scorza 1996; Baggett, Baggett & Anderson 1998), also often associated with barred systems (e.g. van den Bosch & Emsellem 1998). Central velocity-dispersion minima have also been observed in barred galaxies (e.g. Emsellem et al. 2001) and interpreted as the signature of a cold stellar disc, presumably resulting from recent (bar-driven) gas accumulation at the centre (Emsellem et al. 2001; Wozniak et al. 2003). The very weak dependence of rotation on galactic height shown in Fig. 4 is also consistent with NGC 7332 being barred (e.g. Combes et al. 1990; Athanassoula & Misiriotis 2002), although axisymmetric configurations can also in principle give rise to cylindrical rotation (e.g. Rowley 1988).

The only major difference with the kinematic bar signatures emphasized by Bureau & Athanassoula (2004) from  $N$ -body simulations is the strong  $h_3 - V$  anticorrelation in the central parts ( $r \lesssim 7$  arcsec). This is, however, commonly observed in boxy bulges otherwise showing strong evidence for the presence of a bar (Chung & Bureau 2004) and most likely results from the absence of a dissipative component in the simulations (see Friedli & Benz 1993; Wozniak et al. 2003).

The rather homogeneous stellar population of NGC 7332 is also consistent with the presence of a bar. Indeed, the so-called bulge and disc are then both made up of the same (disc) material, and any population gradient that may have been present before the bar formed or may develop afterwards will be smoothed out by the large radial and vertical motions of the stars, at least within the bar region (e.g. Friedli, Benz & Kennicutt 1994; Martin & Roy 1994). Gradients in the very centre may, however, survive or develop due to bar-driven inflows.

The presence of a bar in NGC 7332 was already hinted at by Fisher et al. (1994), who were, however, left to speculate about its existence. Lütticke et al. (2000) estimated a diameter of 56 arcsec for the (weak) bar in NGC 7332 via the characteristic major-axis plateau (observed here in the near-infrared). Considering the discussion above, we can now assert its presence with more confidence. For the remainder of this discussion, we will thus take for granted the presence of a bar in NGC 7332 and focus our efforts on understanding its origin and role in the evolution of the galaxy.

### 7.2 Origin and status of the gas

NGC 7332 exhibits a certain amount of counter-rotating ionized gas ( $\approx 1.5 \times 10^5 M_{\odot}$ ), which has commonly been used to argue for an external origin. In the case of NGC 7332, the ionized gas most likely comes from a tidal interaction with the neighbouring galaxy NGC 7339. We note that extended counter-rotating ionized gas is common in S0s (Bertola et al. 1992), and more specifically in galaxies with boxy bulges (Bureau & Chung in preparation).

Although the ionized gas distribution and kinematics obtained from the SAURON data cube are complex, some of the counter-rotating gaseous component in NGC 7332 seems to have a relatively ordered flow and extends relatively far from the major axis of the galaxy. If we assume that the galaxy is sufficiently inclined, the ionized gas could trace a bar or spiral-like structure in the equatorial plane. If quasi-edge-on, however, it must have a significant vertical extent. Considering the orbital time-scales involved (e.g. about  $2.5 \times 10^6$  yr for a gas clump 1 kpc from the centre), we would

naively expect the (dissipative) counter-rotating ionized gas to be quickly driven back to the equatorial plane. This need not be so, however, as in the tumbling potential of a bar there exists one major family of stable retrograde orbits (the ‘anomalous’ orbits) which is tilted with respect to the equatorial plane and can reach significant heights (Pfenniger & Friedli 1991; Friedli & Udry 1993; Emsellem & Arsenault 1997).

Despite the above ambiguity, it is safe to assume that an accretion event is at the origin of the observed counter-rotating gaseous component in NGC 7332. The same accretion event could be responsible for the formation of the central KDC revealed by SAURON and STIS, as well as the dip in the velocity-dispersion profile (see Section 4.2) and the increase of the Mg *b* and Fe5270 indices at the very centre (see Section 6). The H $\beta$  index values would then set up an upper limit of about 3–4 Gyrs (luminosity-weighted) for the KDC (see Fig. 12). Another later accretion event may be required to explain the more perturbed gas clumps located further away from the equatorial plane (see Section 5), which have rather high relative velocities. This gas has certainly not settled yet, but it will eventually fall back towards the equatorial plane of NGC 7332.

### 7.3 Evolutionary scenarios

Any formation and evolution scenario for NGC 7332 must take into account the two main ingredients hinted at from the observations described in this paper: the presence of a bar and a recent interaction event. We now examine them in turn.

#### 7.3.1 *The role of the bar and the central stellar disc*

In our view, the main morphological and kinematic features that we associate with the bar in Section 7.1 do not represent a proper illustration of bar-driven evolution processes. This is because, although those properties become more extreme as the bar strengthens, most of them are established on time-scales equivalent to the bar formation itself (e.g. Combes et al. 1990; Raha et al. 1991; Bureau & Athanassoula 2004). In the following paragraphs, we thus concentrate on (presumably) bar-driven processes which take place gradually over a much longer time-scale (i.e. tens of dynamical times).

In the context of the bar hypothesis, the interaction between the natural frequencies of the system (e.g. the angular and epicyclic frequencies of the circular orbits) and the pattern speed of the bar implies the existence of resonances, around which the galaxy is shaping. The photometric and kinematic features we observe in NGC 7332 will thus likely be associated with those resonances. The double-disc structure is particularly interesting in this respect since it likely builds up gradually through bar-driven processes (van den Bosch & Emsellem 1998).

The inner disc is usually interpreted as marking the region within the inner Lindblad resonance (ILR). The major-axis plateau observed by Lütticke et al. (2000) extends to a radius of about 25 arcsec ( $\approx 2.8$  kpc), setting a lower limit for the length of the bar. Assuming the standard corotation radius to bar length ratio  $r_{\text{CR}}/r_{\text{b}} = 1.2$  for early-type barred galaxies (e.g. Athanassoula 1992; Gerssen, Kuijken & Merrifield 2003 and references therein),  $r_{\text{CR}} \approx 30$  arcsec. A simple model then predicts the ILR to be at roughly 12 arcsec (see van den Bosch & Emsellem 1998), indeed marking the transition between the inner disc and the outer disc. This may be supported by the line-strength index maps, where there is evidence for a weak decrease of the Fe5015, Mg *b* and Fe5270 profiles at about 10 arcsec along the major axis (see Figs 10 and 11).

A comparison with the edge-on S0 galaxy NGC 4570 appears appropriate here, since it may be at a different stage of a similar (bar-driven) process. NGC 4570 was studied in detail by (van den Bosch & Emsellem (1998), who identified similar morphological features (i.e. double-disc and intermediate boxy structures) to those in NGC 7332. Their study also revealed the same trends in ellipticity and boxiness ( $c_4$ ). The case for a bar was made there almost exclusively based on the identification of two edge-on rings, exceptionally well characterized and consistent with the locations of bar resonances. The structures observed in NGC 4570 are, however, one order of magnitude smaller than in NGC 7332: the ‘nuclear’ disc is less than 1 arcsec in radius and the inner ring is at a radius of  $\approx 1.7$  arcsec. The main difference between the two galaxies then lies in the detection of a large-scale bar of about 3 kpc in NGC 7332, whereas van den Bosch & Emsellem (1998) hinted at the presence of an ‘inner’ bar of  $\approx 320$  pc in NGC 4570 (with no evidence of a large-scale bar). Another obvious qualitative difference is the presence of a significant amount of counter-rotating stars and gas in the central region of NGC 7332. We could speculate on the existence of a secondary bar in NGC 7332, but its detection would require high spatial resolution photometry with high signal-to-noise ratio [e.g. *HST* WFPC-2 or Advanced Camera for Surveys (ACS) images]. Thus, it may well be that NGC 7332 is simply in an earlier stage of a gradual bar-driven evolutionary process.

#### 7.3.2 *The role of interactions*

The formation of decoupled cores is often mentioned within the hierarchical structure formation framework (e.g. Kauffmann, Guiderdoni & White 1994), involving multiple mergers to form galaxies. As mentioned above, although there is no evidence for a recent major merger, the gas distribution and kinematics strongly suggest that NGC 7332 was recently involved in an interaction event (most likely with NGC 7339). The KDC could then correspond to the last or one of the last major accretion (and star-formation) events in such a hierarchy (e.g. Davies et al. 2001). In fact, the bar itself could be the result of that same event, as bars are easily excited by tidal interactions (e.g. Noguchi 1987; Gerin, Combes & Athanassoula 1990).

The neighbour galaxy NGC 7339, a Sb galaxy placed 5 arcmin away from NGC 7332, is most likely the one responsible for such an event. The position of this galaxy is such that the off-plane gas clump found in NGC 7332 could well be a direct continuation of the west side of the disc in NGC 7339. Additionally, the maximum receding velocity measured on the west side of this galaxy is  $\sim 1525$  km s $^{-1}$  (Courteau 1997), which is strikingly similar to the radial velocity of the high-velocity clump in NGC 7332 ( $\sim 1550$  km s $^{-1}$ ). This picture leads to the suggestion that this clump is linked to NGC 7339. Deep H I observations could confirm this idea.

It is hard to assess which of the bar or the interaction occurred first, and whether the KDC was formed from material funnelled by the former or the latter. NGC 7332 will, however, continue to evolve on a time-scale of a few  $10^8$  yr. As emphasized above, the gas will settle down due to its dissipative nature, and some of it may feed the inner few arcseconds (with or without more star formation). In this context, NGC 7332 may be seen as a precursor of the now gas-poor edge-on S0 galaxy NGC 4570.

## 8 CONCLUSIONS

We presented in this paper new SAURON integral-field spectroscopy observations of the edge-on S0 galaxy NGC 7332, along with a discussion of existing ground- and space-based imaging.

The photometric analysis reveals a boxy bulge, a central disc and evidence for the presence of a bar (Section 3; but see also Lütticke et al. 2000). The SAURON observations provide unprecedented coverage of the stellar and gas kinematics. In particular, the stellar kinematics displays a rather smooth velocity field with rotation along the major axis and a weak dependence of rotation on galactic height. We also discovered a stellar kinematically decoupled component (KDC) misaligned with respect to the kinematic major axis of the galaxy, which may be related to a dip in the centre of the velocity-dispersion map ( $r \lesssim 2.5$  arcsec). We finally provided kinematic evidence for the presence of a larger central stellar disc ( $r \lesssim 7$  arcsec).

NGC 7332 presents significant ionized gas emission in the spectral range studied, in particular in the  $H\beta$  (4861 Å) and [O III] (4959 and 5007 Å) lines. We used a sophisticated technique to separate the absorption and emission lines, yielding both pure absorption spectra from which to derive the stellar kinematics and pure emission spectra for the gas kinematics (and distribution). The emission maps reveal a very complex gas morphology and kinematics. This is found especially in [O III], which is mainly counter-rotating with respect to the stars. The gas maps (Fig. 9) show significant amounts of gas at high relative velocities outside of the equatorial plane, whereas the central parts display rather ordered motions. We also found traces of a faint corotating ionized gas component.

The analysis of the absorption line-strengths in NGC 7332 show that its stellar populations are generally young ( $5 \pm 2$  Gyr), not only in the disc but also in the bulge, in agreement with previous studies (Balcells & Peletier 1994; Vazdekis et al. 1996; Terlevich & Forbes 2002). The metallicity indices (i.e. Fe5015, Mgb, Fe5270) show an increase in the centre, contrasting with the rather homogeneous  $H\beta$  index. There is also weak evidence of a decrease at  $r \approx 10$  arcsec, between the central and the outer discs.

The existence of a large-scale bar in NGC 7332 can simultaneously explain most of the features found in the galaxy. The boxy morphology can be explained by the thickness of the bar after buckling, which also leads to homogeneous stellar populations across the bulge and disc. The stellar kinematics can similarly be explained by the particular orbital structure of barred discs (see Bureau & Athanassoula 2004). The KDC, on the other hand, was most likely formed through late gas infall, an accretion event perhaps related to the complex ionized gas distribution currently seen.

As emphasized by the unique data set presented in this paper, the S0 galaxy NGC 7332 possesses a number of peculiar properties helping to unravel its formation and evolution processes. There is strong observational evidence that interactions played a role in the shaping of NGC 7332, while the bar has and is certainly significantly affecting its evolution. Given the amount of gas yet to settle and the relatively young age of the stellar populations in NGC 7332, it probably represents an earlier evolutionary stage of more regular edge-on S0 galaxies (i.e. NGC 4570, van den Bosch & Emsellem 1998; NGC 3115, Emsellem 2003; NGC 3377, Bacon et al. 2001). The morphology and dynamics of present-day S0 galaxies can thus probably be understood through the combined effects of both interactions and bar-driven processes.

FITS files with the kinematical and line-strength maps presented in this paper are available on request from the first author.

## ACKNOWLEDGMENTS

We would like to thank Fabien Wernli for stimulating discussions in the first stages of this work. The William Herschel Telescope is operated on the island of La Palma by the Isaac Newton Group in

the Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. Support for this work was provided by NASA through Hubble Fellowship grant *HST*-HF-01136.01 awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS 5-26555. MB is a Hubble Fellow and MC acknowledges support from a VENI grant awarded by the Netherlands Organization of Scientific Research (NWO).

## REFERENCES

- Andredakis Y. C., Peletier R. F., Balcells M., 1995, *MNRAS*, 275, 874  
 Athanassoula E., 1992, *MNRAS*, 259, 345  
 Athanassoula E., Misiriotis A., 2002, *MNRAS*, 330, 35  
 Bacon R. et al., 2001, *MNRAS*, 326, 23  
 Baggett W. E., Baggett S. M., Anderson K. S. J., 1998, *AJ*, 116, 1626  
 Balcells M., Peletier R. F., 1994, *AJ*, 107, 135  
 Bender R., 1988, *A&A*, 202, L5  
 Bertola F., Buson L. M., Zeilinger W. W., 1992, *ApJ*, 401, L79  
 Bureau M., Athanassoula E., 2004, *ApJ*, submitted  
 Bureau M., Freeman K. C., 1999, *AJ*, 118, 126  
 Caldwell N., 1983, *ApJ*, 268, 90  
 Cappellari M., Copin Y., 2003, *MNRAS*, 342, 345  
 Cappellari M., Emsellem E., 2004, *PASP*, 116, 138  
 Cappellari M., Verolme E. K., van der Marel R. P., Verdoes G. A. V., Illingworth G. D., Franx M., Carollo C. M., de Zeeuw P. T., 2002, *ApJ*, 578, 787  
 Cardiel N., 1999, PhD thesis, Univ. Complutense de Madrid  
 Carollo C. M., 1999, *ApJ*, 523, 566  
 Carter D., 1978, *MNRAS*, 182, 797  
 Chung A., Bureau M., 2004, *AJ*, in press  
 Combes F., Debbasch F., Friedli D., Pfnegner D., 1990, *A&A*, 233, 82  
 Courteau S., 1997, *AJ*, 114, 2404  
 Courteau S., de Jong R. S., Broeils A. H., 1996, *ApJ*, 457, L73  
 Davies R. L. et al., 2001, *ApJ*, 548, L33  
 de Vaucouleurs G., 1959, *Handbuch der Physik*, 53, 275  
 de Vaucouleurs G., de Vaucouleurs A., Corwin H. G., Buta R. J., Paturel G., Fouque P., 1991, *Sky & Telescope*, 82, 621  
 de Zeeuw P. T. et al., 2002, *MNRAS*, 329, 513  
 Dressler A., 1980, *ApJ*, 236, 351  
 Emsellem E., 2003, in Ho L. C., ed., *Carnegie Observatories Astrophysics Series, Vol. 1, Coevolution of Black Hole and Galaxies*. Cambridge Univ. Press, Cambridge (available online at <http://www.ociw.edu/ociw/symposia/series/symposium1/proceedings.html>)  
 Emsellem E., Arsenault R., 1997, *A&A*, 318, L39  
 Emsellem E., Greusard D., Combes F., Friedli D., Leon S., Pécontal E., Wozniak H., 2001, *A&A*, 368, 52  
 Faber S. M., Gallagher J. S., 1976, *ApJ*, 204, 365  
 Fisher D., 1997, *AJ*, 113, 950  
 Fisher D., Illingworth G., Franx M., 1994, *AJ*, 107, 160  
 Franx M., Illingworth G. D., 1988, *ApJ*, 327, L55  
 Freeman K. C., 1970, *ApJ*, 160, 811  
 Friedli D., Benz W., 1993, *A&A*, 268, 65  
 Friedli D., Udry S., 1993, in DeJonghe H., Habing H. J., eds, *Proc. IAU Symp. Vol. 153, Galactic Bulges*. Kluwer, Dordrecht, p. 273  
 Friedli D., Benz W., Kennicutt R., 1994, *ApJ*, 430, L105  
 Gerin M., Combes F., Athanassoula E., 1990, *A&A*, 320, 37  
 Gerssen J., Kuijken K., Merrifield M. R., 2003, *MNRAS*, 345, 261  
 González J. J., 1993, PhD thesis, Univ. California, Santa Cruz  
 Goudfrooij P., Baum S. A., Walsh J. R., 1997, *BAAS*, 29, 1239  
 Gregg M., 1989, *ApJ*, 337, 45  
 Gunn J. E., Gott J. R. I., 1972, *ApJ*, 176, 1  
 Hubble E. P., 1936, *The Realm of the Nebulae*. Yale Univ. Press, New Haven  
 Kauffmann G., Guiderdoni B., White S. D. M., 1994, *MNRAS*, 267, 981  
 Kuntschner H., 2000, *MNRAS*, 315, 184  
 Kuntschner H., Davies R. L., 1998, *MNRAS*, 295, L29

- Kuntschner H., Lucey J. R., Smith R. J., Hudson M. J., Davies R. L., 2001, MNRAS, 323, 615
- Larson R. B., Tinsley B. M., Caldwell C. N., 1980, ApJ, 237, 692
- Lauer T. R. et. al., 1995, AJ, 110, 2622
- Lütticke R., Dettmar R.-J., Pohlen M., 2000, A&A, 362, 435
- Martin P., Roy J.-R., 1994, ApJ, 424, 599
- Merrifield M. R., Kuijken K., 1999, A&A, 345, L47
- Noguchi M., 1987, MNRAS, 228, 635
- Osterbrock D. E., 1989, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei*. Univ. Science Books, Mill Valley, CA, p. 422
- Peletier R. F., Balcells M., 1996, AJ, 111, 2238
- Peletier R. F., Balcells M., 1997, *New Astron.*, 1, 349
- Peletier R. F., Balcells M., Davies R. L., Andredakis Y., Vazdekis A., Burkert A., Prada F., 1999, MNRAS, 310, 703
- Pfenniger D., 1993, DeJonghe H., Habing H. J., eds, *Proc. IAU Symp. Vol. 153, Galactic Bulges*. Kluwer, Dordrecht, p. 387
- Pfenniger D., Friedli D., 1991, A&A, 252, 75
- Plana H., Boulesteix J., 1996, A&A, 307, 391
- Raha N., Sellwood J. A., James R. A., Kahn F. D., 1991, Nat, 352, 411
- Rowley G., 1988, ApJ, 331, 124
- Sandage A., 1961, *The Hubble Atlas of Galaxies*. Carnegie Institution, Washington
- Sandage A., Visvanathan N., 1978, ApJ, 225, 742
- Schweizer F., 1986, Sci, 231, 227
- Seifert W., Scorza C., 1996, A&A, 310, 75
- Sellwood J. A., Wilkinson A., 1993, *Rep. Prog. Phys.*, 56, 173
- Statler T. S., Smecker-Hane T., 1999, AJ, 117, 839
- Terlevich A. I., Forbes D. A., 2002, MNRAS, 330, 547
- Terndrup D. M., Davies R. L., Frogel J. A., Depoy D. L., Wells L. A., 1994, ApJ, 432, 518
- Thomas D., Maraston C., Bender R., 2003, MNRAS, 339, 897
- Tonry J. L., Dressler A., Blakeslee J. P., Ajhar E. A., Fletcher A., Luppino G. A., Metzger M. R., Moore C. B., 2001, ApJ, 546, 681
- Toomre A., 1977, *ARA&A*, 15, 437
- Trager S. C., Faber S. M., Worthey G., González J. J., 2000, AJ, 119, 1645
- van den Bergh S., 1960a, ApJ, 131, 558
- van den Bergh S., 1960b, ApJ, 131, 215
- van den Bosch F. C., Emsellem E., 1998, MNRAS, 298, 267
- van der Marel R. P., Franx M., 1993, ApJ, 407, 525
- Vazdekis A., 1999, ApJ, 513, 224
- Vazdekis A., Arimoto N., 1999, ApJ, 525, 144
- Vazdekis A., Casuso E., Peletier R. F., Beckman J. E., 1996, ApJS, 106, 307
- Wagner S. J., Bender R., Moellenhoff C., 1988, A&A, 195, L5
- Wernli F., Emsellem E., Copin Y., 2002, A&A, 396, 73
- White S. D. M., Rees M. J., 1978, MNRAS, 183, 341
- Worthey G., Faber S. M., Gonzalez J. J., Burstein D., 1994, ApJS, 94, 687
- Worthey G., Ottaviani D. L., 1997, ApJS, 111, 377
- Wozniak H., Combes F., Emsellem E., Friedli D., 2003, A&A, 409, 469

This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.