

**Parasitic Light in NGST instruments:
effects of filter leaks in the visible and near IR camera.**

Stefano Cristiani

*ST European Coordinating Facility, European Southern Observatory
Karl-Schwarzschild-Strasse 2, D-85748 Garching bei Muenchen*

Stephane Arnouts

*European Southern Observatory
Karl-Schwarzschild-Strasse 2, D-85748 Garching bei Muenchen*

Robert A.E. Fosbury

*ST European Coordinating Facility, European Southern Observatory
Karl-Schwarzschild-Strasse 2, D-85748 Garching bei Muenchen*

ABSTRACT

A detailed analysis of NGST multiband photometry applied to the reference case of the study of high-redshift galaxies has been carried out with simulations based on galaxy SEDs derived from the currently available empirical and model templates and on plausible standard filter-sets.

In order to correctly identify star forming galaxies in the redshift range $5 < z < 9$ and early-type galaxies above $z > 2$ and avoid confusion with other SEDs it is mandatory to have photometric information in optical bands, besides a standard IR filter-set like *F110 F160 K L M*. In particular by adding the *V606, I814* and *Z – Gunn* filters a good discrimination is obtained above $z > 5$ for star forming galaxies and $z > 1$ for early-types. The case for an extension of the NGST wavelength domain to the optical range is therefore strongly supported by this analysis.

The effects of leaks in the filter blocking have also been investigated. In spite of rather pessimistic assumptions (a constant leak at a level of 10^{-4} of the peak transmission over the whole spectral range or a leak of Gaussian shape placed at 1.5 times the effective wavelength of the filter with an amplitude of 10^{-3} of the peak transmission and a width of 5% of the central wavelength) the effects are not dramatic: the accuracy of the determination of photometric redshifts with a standard *V606 I814 Z – Gunn F110 F160 K L M* filter set is not significantly deteriorated for a sample limited to $M_{AB} = 28$. In the range $5.5 < z < 12.5$ a rather good accuracy a typical error $|z_{sim} - z_{phot}| \leq 0.2$ is achieved.

With the filter blocking achieved by the present standard technology, extended and continuous spectral coverage appears to be the driving factor to maximize the scientific output.

1. Introduction

The NGST Design Reference Mission (DRM) is a representative science program elaborated in sufficient detail to aid in the development of functional requirements for the NGST mission. One of the main results of the DRM has been the identification of a large fraction of subprograms (8 out of 12) requiring observations at optical wavelengths. Such an extension poses significant technical challenges. In particular it requires an efficient blocking of photons at undesired wavelengths over a range covering more than three octaves. In the following we examine the consequences of leaks in the transmission of filters for a typical programme of determination of photometric redshifts of high- redshift galaxies.

2. Baseline of the Instrument

Table 1 shows the main characteristics of the Visible-NIR camera as conceived by the Visible and NIR camera Subcommittee of the “Ad Hoc Science Working Group” (ASWG). Two options are envisaged: an *optimal* and a *bare bones* version. More details can be found at the URL <http://www.stecf.org/ngst/instruments.html>.

3. Band-passes of the filters

We have assumed as reference bandpasses for the filters of the optical-NIR camera the standard *V606 I814 Z–Gunn F110 F160 K L M* set commonly used on HST with WFPC2 (*V606 I814*) or NICMOS (*F110 F160*) and from the ground (Fig. 1). Details about the filter properties are given in Table 2

The actual choice of the filters to be used in space with NGST may be different but the arguments developed in the following do not depend critically on this detail.

Table 1. Visible-NIR Camera

	Optimal	Bare Bones
λ range	0.6 – 5 μ m	1.0 – 5.0 μ m
Sampling	0.03"	0.03"
FOV	4' \times 4'	2' \times 2'
$\lambda/\Delta\lambda$	≤ 10	≤ 10

Table 2. Properties of the assumed filter set

Name	$\langle \lambda \rangle$ (\AA)	λ_{eff} (\AA)	FWHM (\AA)	M_{AB} (Vega)
F606	6031.0	5831.6	2050	0.096
F814	8011.4	7862.3	1400	0.417
Zgunn	9124.8	9066.8	1200	0.527
F110	11287.8	10493.9	5750	0.698
F160	16095.6	15798.5	3750	1.313
Ks	21639.7	21522.1	2750	1.841
L	37788.6	37497.2	5800	2.917
M	47656.8	47374.8	6400	3.380




Fig. 1.— The reference filter set used for the simulations.

4. Typical SEDs of the sources and the locus of galaxies in multicolor space

To simulate a range of galaxy SEDs we have adopted the templates of Coleman et al. (1980) for a typical elliptical, Sbc, Scd and Irregular galaxy plus a GISSEL (Bruzual and Charlot 2001) template for a very blue galaxy (with a constant star formation rate and an age of 0.1 Gyr). The SEDs are shown in Fig. 2 and in the following will be referred to as extended CWW templates.

Fig. 3 shows the evolution of the apparent magnitude of typical galaxies of $10^{11}M_{\odot}$ in the $F606$ and L bands and gives a rough idea of the magnitudes of interest. It can be appreciated how an Elliptical fades away relatively quickly in the $F606$ band, while other star-forming galaxies remain - as observed - significantly brighter.

The upper panel of Fig. 4 shows the evolution of the colors of the galaxies described in the previous Section in a $(F606 - F814)$ vs. $(F110 - K)$ plane. The color tracks are limited to $z < 3.5$ for an Elliptical SED and to $z < 5$ for Sbc galaxies. With this combination of bandpasses it is straightforward to identify star-forming galaxies with $z > 5$ and Ellipticals at $z > 1$. The lower panel shows the same in a $(F110 - F160)$ vs. $(F160 - K)$ plane and demonstrates how without optical bands it becomes difficult to isolate star-forming galaxies at $z < 9$.

5. Photometric redshifts and the effects of “(red) leaks”

In order to quantify the different performances of different sets of filters and the effects of “red-leaks” in a filter configuration extending from the visible to the infrared we have carried out simulations of the assignment of photometric redshifts to a sample of faint galaxies. To this end, we have tried to generate a simulated photometric catalog covering a very large range of galaxy spectral properties and to estimate photometric redshifts from it using an economic as independent as possible set of templates. Galaxies were simulated in a redshift interval $0 < z < 20$ according to GISSEL library. This spectral synthesis model is controlled by a number of free parameter. The star formation rate for a galaxy at a given age is governed by the assumed $e - folding$ star formation time-scale τ . Several values of τ and galaxy ages are necessary to reproduce the different observed spectral types. For the shape of the initial mass function (IMF) we have assumed a Salpeter law, which turns out not to be a critical choice. In addition to the GISSEL parameterization, we have added the internal reddening of each galaxy by applying the observed attenuation law for local starburst galaxies by Calzetti et al. (1997), with the E_{B-V} normalization varying between 0 and 0.2 mag. We have also included the Lyman absorption produced by the intergalactic medium as a function of redshift. More details on this type of simulations can be found in Arnouts et al. (1999). Fig. 5 shows the galaxy counts as a function of the apparent magnitude, the age and redshift distributions, the $z - mag$ plane characterizing the simulation.

Magnitudes and colors have been derived from the simulated SEDs by convolution with the filter passbands described in Sect.3. Measurement errors have been added, according to the NGST mission simulator (<http://www.ngst.stsci.edu/nms/main/>),




Fig. 2.— The spectral energy distribution of five template galaxies.





Fig. 3.— Evolution of the apparent magnitude of typical galaxies of $10^{11}M_{\odot}$.




Fig. 4.— Redshift evolution of the colors of galaxies. Upper panel: $(F606 - F818)$ vs. $(F110 - K)$. Lower panel: $(F110 - F160)$ vs. $(F160 - K)$.




Fig. 5.— Characteristic distributions of the simulation of galaxies derived from GISSEL models: counts as a function of the apparent magnitude (upper left), redshift distribution (upper and lower right), the $z - mag$ plane (lower left). In the first two panels the dashed lines refer to galaxies with ages larger than 1 Gyr, continuous lines to ages between 10^8 and 10^9 yr.

assuming an exposure time of 10^4 s.

From this set of “simulated observed colors” photometric redshifts have been derived using a standard procedure of χ^2 optimization, comparing the observed fluxes with errors to the extended CWW set of templates described in Sect. 4.

The χ^2 has been computed as:

$$\chi^2 = \sum_i \left[\frac{F_{observed,i} - A \cdot F_{template,i}}{\sigma_i} \right]^2 \quad (1)$$

where $F_{observed,i}$ is the flux observed in a given filter i , σ_i is its uncertainty, $F_{template,i}$ is the flux of the template in the same filter and the sum is over the various filters. The template fluxes have been normalized to the observed ones with the factor A minimising the χ^2 value :

$$s = \sum_i \left[\frac{F_{observed,i} \cdot F_{template,i}}{\sigma_i^2} \right] / \sum_i \left[\frac{F_{template,i}^2}{\sigma_i^2} \right] \quad (2)$$

Fig. 6 shows the comparison between photometric redshift estimates obtained with a *V606 I814 Z – Gunn F110 F160 K L M* filter set and a filter set limited to the infrared bands. As suggested in the previous section in order to correctly identify star forming galaxies in the redshift range $5 < z < 9$ and avoid confusion with other SEDs it appears mandatory to have, besides the IR filters, photometric information in the optical bands. In the following we will adopt the set *V606 I814 Z – Gunn F110 F160 K L M* as standard.

In order to explore the effects of filter leaks the following approach was adopted:

1. two types of leaks have been foreseen: a constant leak at a level of 10^{-4} of the peak transmission over the whole spectral range and a leak of Gaussian shape placed at 1.5 times the effective wavelength of the filter with an amplitude of 10^{-3} of the peak transmission and a width of 5% of the central wavelength. The effect of Gaussian leaks on the filter transmission is shown in Fig. 9
2. The leaks have been included in the system spectral response and new magnitudes and colors have been produced with the GISSEL models.
3. The new colors have been used to produce photometric redshifts with the extended CWW templates. No knowledge of the leaks was used in this step, i.e. the system responses assumed in the z -phot estimation are those of Fig. 1, without leaks.

The typical errors deriving from photometric uncertainties and unaccounted leaks are illustrated in Fig. 7-8 and 10-11. The dot-dashed, long-dashed, short-dashed and continuous lines correspond to the 95, 90, 68 and 50 percentile of the absolute deviation $|z_{sim} - z_{phot}|$, respectively. It can be seen that errors are in all cases relatively large at $z \leq 5$. This is an obvious effect due to the lack of bandpasses bluer than V_{606} , and the consequent difficulty to identify the Lyman break at relatively low redshift and distinguish it, for example, from the 4000 Å break at lower redshift.




Fig. 6.— Comparison between the accuracy of the photometric redshift estimates with a filter set including optical bands (upper panel) and a filter set limited to infrared bands (lower panel). The dot-dashed, long-dashed, short-dashed and continuous lines correspond to the 99, 90, 68 and 50 percentile of the absolute deviation $|z_{sim} - z_{phot}|$, respectively. The galaxy sample is limited to $M_{AB} < 28$




Fig. 7.— Typical errors of the photometric redshifts with a $V606 I814 Z - Gunn F110 F160 K L M$ filter set. The upper panel shows the result obtained assuming only the standard photometric error, the lower panel includes the effects of a constant leak at a level of 10^{-4} of the peak transmission over the whole spectral range. The dot-dashed, long-dashed, short-dashed and continuous lines correspond to the 95, 90, 68 and 50 percentile of the absolute deviation $|z_{sim} - z_{phot}|$, respectively. The galaxy sample is limited to $M_{AB} < 27$.




Fig. 8.— Same as Fig. 7, but down to $M_{AB} = 28$.




Fig. 9.— The response of the *V606 I814 Z – Gunn F110 F160 K L M* filter with the addition of a leak of Gaussian shape placed at 1.5 times the effective wavelength of the filter with an amplitude of 10^{-3} of the peak transmission and a width of 5% of the central wavelength.




Fig. 10.— Typical errors of the photometric redshifts with a $V606 I814 Z - Gunn F110 F160 K L M$ filter set. The upper panel shows the result obtained assuming only the standard photometric error, the lower panel includes the effects of a leak of Gaussian shape placed at 1.5 times the effective wavelength of the filter with an amplitude of 10^{-3} of the peak transmission and a width of 5% of the central wavelength. The dot-dashed, long-dashed, short-dashed and continuous lines correspond to the 95, 90, 68 and 50 percentile of the absolute deviation $|z_{sim} - z_{phot}|$, respectively. The galaxy sample is limited to $M_{AB} < 27$.




Fig. 11.— Same as Fig. 10 but down to $M_{AB} = 28$.

The effect of the leaks, in spite of the rather pessimistic assumptions, is not dramatic. We observe in general a moderate worsening of the error level, but there is no qualitative change of the accuracy. No relevant difference is found between the continuous and the Gaussian leak cases.

In the range $5.5 < z < 12.5$ a rather good accuracy is achieved with a typical error $|z_{sim} - z_{phot}| \leq 0.2$ in all cases down to $M_{AB} = 28$. The situation deteriorates slightly around $z \simeq 15$, but the result is due mainly to the insufficient sampling of the SEDs between the K_s and the M filters, in particular to the “hole” at $2.9 \mu\text{m}$.

6. Conclusions

A detailed analysis of NGST multiband photometry applied to the reference case of the study of high-redshift galaxies has been carried out with simulations based on galaxy SEDs derived from the currently available empirical and model templates and on plausible standard filter-sets.

In order to correctly identify star forming galaxies in the redshift range $5 < z < 9$ and early-type galaxies above $z > 2$ and avoid confusion with other SEDs it is mandatory to have photometric information in optical bands, besides a standard IR set like $F110 F160 K L M$. In particular by adding the $V606$, $I814$ and Z – Gunn filters a good discrimination is obtained above $z > 5$ for star forming galaxies and $z > 1$ for early-types. The case for an extension of the NGST wavelength domain to the optical range is therefore strongly supported by this analysis.

The effects of leaks in the filter blocking have also been investigated. In spite of rather pessimistic assumptions (a constant leak at a level of 10^{-4} of the peak transmission over the whole spectral range or a leak of Gaussian shape placed at 1.5 times the effective wavelength of the filter with an amplitude of 10^{-3} of the peak transmission and a width of 5% of the central wavelength) the effects are not dramatic: the accuracy of the determination of photometric redshifts with a standard $V606 I814 Z$ – Gunn $F110 F160 K L M$ filter set is not significantly deteriorated for a sample limited to $M_{AB} = 28$. In the range $5.5 < z < 12.5$ a rather good accuracy a typical error $|z_{sim} - z_{phot}| \leq 0.2$ is achieved.

With the filter blocking achieved by standard technology at present, extended and continuous spectral coverage appears to be the driving factor to maximize the scientific output.

7. Acknowledgments

We are grateful to A.Moorwood for enlightening discussions about the performances of IR filters and detectors.

REFERENCES

- Arnouts S., Cristiani S., Moscardini L., Matarrese S., Lucchin F., Fontana A., Giallongo E., 1999, MNRAS, 310, 540
- Bruzual, G., Charlot, S., 2001 (GISSEL) in preparation
- Calzetti D., 1997, in Waller W.H. et al. eds., The Ultraviolet Universe at Low and High Redshift: Probing the Progress of Galaxy Evolution, AIP Conf. Proc. 408. AIP, Woodbury, 403
- Coleman, G.D., Wu, C.C., Weedman, D.W. 1980, ApJS, 43, 393 (CWW)