

The SPHERE view of the Taurus star-forming region

The full census of planet-forming disks with GTO and DESTINYs programs

A. Garufi¹, C. Ginski², R. G. van Holstein³, M. Benisty^{4,5}, C. F. Manara⁶, S. Pérez^{7,8}, P. Pinilla⁹, A. Ribas¹⁰,
P. Weber^{7,8}, J. Williams¹¹, L. Cieza^{12,7}, C. Dominik¹³, S. Facchini¹⁴, J. Huang¹⁵, A. Zurlo^{12,7}, J. Bae¹⁶,
J. Hagelberg¹⁷, Th. Henning¹⁸, M. R. Hogerheijde^{19,13}, M. Janson²⁰, F. Ménard⁵, S. Messina²¹,
M. R. Meyer²², C. Pinte^{23,5}, S. Quanz²⁴, E. Rigliaco²⁵, V. Roccatagliata¹, H. M. Schmid²⁴, J. Szulágyi²⁴,
R. van Boekel¹⁸, Z. Wahhaj³, J. Antichi¹, A. Baruffolo²⁵, and T. Moulin⁵

(Affiliations can be found after the references)

Received -; accepted -

ABSTRACT

The sample of planet-forming disks observed by high-contrast imaging campaigns over the last decade is mature enough to enable the demographic analysis of individual star-forming regions. We present the full census of Taurus sources with VLT/SPHERE polarimetric images available. The whole sample sums up to 43 targets (of which 31 have not been previously published) corresponding to one fifth of the Class II population in Taurus and about half of the observable such objects. A large fraction of the sample are apparently isolated, faint disks (equally divided between small and large, self-shadowed disks). Ambient signal is visible in about one third of the sample. This probes the interaction with the environment and with companions or the outflow activity of the system. The central portion of the Taurus region almost exclusively hosts faint disks while the periphery also hosts bright disks interacting with their surroundings. The few bright disks are found around apparently older stars. The overall picture is that the Taurus region is in an early evolutionary stage of planet formation. Yet, some objects are discussed individually as in an intermediate or exceptional stage of the disk evolution. This census provides a first benchmark for the comparison of the disk populations in different star forming regions.

Key words. protoplanetary disks – techniques: polarimetric – star-forming regions: Taurus

1. Introduction

The study of the planet formation is hampered by the paucity of detectable protoplanets in their natal disks. The other most direct vehicle to determine observationally how planets form are resolved maps of the planet-forming disk itself, and in particular of the (evolving) morphological features that are connected to the interaction with embedded planets. This approach has been successfully carried out over the last few years thanks to the near-IR (NIR) high-contrast imaging from the ground (see review by Benisty et al. 2023) and the high-resolution imaging granted by the Atacama Large (sub-)Millimeter Array (ALMA, see the review by Andrews 2020). While the ALMA community has soon carried out demographical studies at moderate resolution of entire star-forming regions (e.g., Pascucci et al. 2016; Ansdell et al. 2016), such an effort by the NIR community struggled to take off mostly because of the longer minimum exposure times required on the telescopes.

About one decade after the onset of the latest-generation instruments offering high-contrast imaging (such as GPI, Macintosh et al. 2014, and SPHERE, Beuzit et al. 2019), the community is finally in the condition to lead demographical studies on large samples of young stellar objects. This effort has been made possible in particular by systematic programs like the SEEDS at Subaru (e.g., Hashimoto et al. 2012; Mayama et al. 2012), the Guaranteed Time Observation (GTO) program at SPHERE (e.g., Garufi et al. 2016; Ginski et al. 2016), DARTTS at SPHERE (Avenhaus et al. 2018; Garufi et al. 2020), LIGHTS at GPI (e.g.,

Laws et al. 2020; Rich et al. 2022), and more recently by DESTINYs at SPHERE (e.g., Ginski et al. 2021, 2022).

In particular, DESTINYs (Disk Evolution Study Through Imaging of Nearby Young Stars) is an ESO large program to obtain deep, polarized intensity images of 85 young stellar objects in all nearby star forming regions with the goal to alleviate the bias on the available sample and constrain the disk evolution over a large range of ages. Thanks to DESTINYs and the previous programs, the demography of disks can now be studied through a meaningfully large and reasonably little biased sample. A first taxonomical analysis of the full sample available at the time (58 targets) was performed by Garufi et al. (2018) who identified six main classes of disk types based on NIR images. More recently, Rich et al. (2022) analyzed the 44 sources in LIGHTS to significantly enlarge the stellar parameter space to higher masses. Furthermore, Garufi et al. (2022) examined a sample of 15 faint disks to demonstrate the high occurrence of self-shadowed disks, where the inner portion of the disk casts a shadow on the outer disk that is observable by the current generation of telescopes.

In this work, we take a step forward in the study of disk evolution by moving our objective from the specific type of objects or observing program to the population of young stars in a specific star-forming region. In parallel with the work by Ginski et al. on Chamaeleon and by Viegas et al. on Orion, we collect the entirety of the SPHERE dataset on the Taurus region that has been taken over the last decade with SPHERE/IRDIS (Dohlen et al. 2008) in Dual-beam Polarimetric Imaging mode (DPI, de Boer et al. 2020; van Holstein et al. 2020) by the SPHERE GTO

(PI: Beuzit), by the large program DESTINYS (PI: Ginski), and by individual open-time programs. The final sample that we build is composed of 43 targets (of which 31 have not been previously published), making it the largest sample of high-contrast images from an individual region ever studied as a whole.

The Taurus-Auriga complex is one of the most studied, nearby (128–198 pc, Galli et al. 2018) low-mass star-forming regions. It hosts a few hundred young stellar objects that are distributed in small kinematical groups (Esplin & Luhman 2019; Joncour et al. 2018). All Taurus members appear coeval (1–3 Myr, Luhman 2023) and yet they are in very diverse evolutionary stages, ranging from embedded protostars to disk-free stars (Kenyon et al. 2008). Large-scale maps of the dust distribution revealed that Taurus lies on the near hemispheric side of the so-called Perseus-Taurus shell (Leike et al. 2020), which suggests that the whole cloud may have been formed via compression of material between the expanding Perseus-Taurus bubble and the local bubble (Soler et al. 2023).

For a long time, the large sky area and the low density of Taurus hindered the determination of the individual membership. The *Gaia* mission (Gaia Collaboration 2016) is now significantly reducing contamination from field stars by providing kinematical data on all stars as faint as 20 mag in the *G* waveband, corresponding to a stellar mass of 0.05 M_{\odot} at the distance of Taurus (Luhman 2023). Several authors have employed the *Gaia* data to constrain the morphological and kinematical properties of Taurus (e.g., Galli et al. 2019; Roccatagliata et al. 2020; Krolikowski et al. 2021). In this work, we make use of the membership catalogue created by Luhman (2023) based on the latest *Gaia* release (DR3, Gaia Collaboration et al. 2022). We focus on the specific evolutionary stage of Class II objects (Lada 1987) as these stars still host a gas-rich planet-forming disk but no longer possess any dusty envelope that obscures the optical light from the star.

The paper is organized as follows. In Sect. 2, we census the spatial distribution of the available SPHERE sample, and calculate the stellar and disk properties. In Sect. 3, we describe the observations and data reduction. The results from the SPHERE images are given in Sect. 4 while those from the comparison with literature ALMA images in Sect. 5. Finally, we discuss and summarize our results in Sects. 6 and 7.

2. Sample

The sample studied in this work consists of the 43 sources in Taurus with available VLT/SPHERE DPI images (see Sect. 3). In this section, we discuss their distribution in space and in stellar and disk properties with the intent to set a background for the SPHERE observations described in Sect. 4.

2.1. Spatial distribution of the sample

Figure 1 shows the spatial distribution of the 43 sources studied in this work. A quick look to their distribution already gives a hint of their membership to different groups. Following the kinematical classification by Luhman (2023) based on *Gaia* DR3 (Gaia Collaboration et al. 2022), our targets belong to 11 different groups and one association near Taurus (namely HD35187 that hosts CQ Tau and MWC758). The groups that are most represented by our sample are L1524/L1529/B215 (10 members) and L1517 (7 members) while in five cases we only have one member. The membership and the distance from *Gaia* for the entire sample are shown in Appendixes A and B. With some notable exceptions (see below), the distance of each source is close

to the averaged distance of the respective group, spanning from the 130 pc of L1495/B209 to the 196 pc of L1558.

The goodness of the *Gaia* astrometric observation used in this work is commonly evaluated through the renormalized unit weight error (RUWE). A single star with a good astrometric solution has a RUWE around 1.0 or anyway smaller than 1.4. Interestingly, only 19 of our 43 sources have RUWE smaller than 1.4, and these are all single stars (see Sect. 2.3). Fitton et al. (2022) proposed to adopt a higher threshold (2.5) to evaluate the goodness of the astrometric solution on disk-bearing stars. With this threshold, 29 of 43 our sources would have a good solution. Eight of our targets have a RUWE larger than 4 meaning that their astrometric solution is too poor to lend credence to the calculated distance. In one of these eight cases, GG Tau A, we use the distance of a nearby, bound companion (GG Tau B) with a good solution. The other seven cases are marked in Table B.1. Interestingly, among these seven cases we find the aforementioned discrepancies between the individual distance and the averaged distance of the respective group (e.g., RW Aur at 183 pc from the group L1517, at 156 pc, or V807 Tau at 184 pc from L1524, lying at 128 pc), pointing to an erroneous parallax solution.

2.2. This sample within the whole Taurus population

Based on *Gaia* DR3, Luhman (2023) identified 532 Taurus members. From the nearly identical sample by Esplin & Luhman (2019), approximately 200 Taurus members have an SED indicative of a full Class II disk around the star, or ~220 including transition and evolved disks (with most of the remaining sources being Class III). Given the sample size of this work, we conclude that about one fifth of the Taurus planet-forming disks have been probed by direct near-IR imaging.

The main technical limit toward the high-contrast imaging of Class II objects within 200 pc is the stellar brightness at optical wavelengths needed to drive the adaptive-optics system at the telescope. For SPHERE, this limit is set to an *R* waveband magnitude of 13 (although an optimal performance is only expected up to 11). From the whole sample by Esplin & Luhman (2019), 187 targets show a *Gaia* G_{RP} passband¹ magnitude smaller than 13. Of these, 82 are classified as either Class II or transition-disk objects. Therefore, about a half² of the planet-forming disks in Taurus that are potentially observable with direct near-IR imaging have been surveyed. However, if we focus on the 49 Class II targets with *Gaia* G_{RP} smaller than 12 (for which the adaptive-optics performance is at worst sub-optimal) then more than 75% (that is 37) of the possible targets have been observed. The distribution of all Taurus sources, of all Class II sources, and of all the observable Class II sources is shown in Appendix A.

The whole Taurus sample is limited in both distance and age. Therefore, the cutoff in *R* magnitude on the available sample reflects in a cutoff on the stellar mass only and can be set to approximately 0.4 M_{\odot} . This value is in fact the lowest limit of our sample (see Sect. 2.3). The only other selection criterion that may hold is an IR excess indicative of a class II object. We can therefore conclude that our sample is unbiased and moderately complete within the Taurus stars more massive than 0.4 M_{\odot} hosting a full or a transition disk. It however remains completely blind

¹ the *Gaia* G_{RP} passband is wider in wavelength than the Johnson *R*. The exact conversion depends on the stellar color but for our exercise is not necessary as the difference is relatively small (1 mag at most).

² $40 \div 82 = 49\%$, where 40 is our sample size (43) minus CQ Tau and MWC758 that are formally not in Taurus (see Sect. 2.1), and V1025 Tau that is classified as a Class III (see Sect. 4.3.2).




Fig. 1. Spatial distribution of the sample. The background image is the Taurus region at 100 μm (Band 4) from the Improved Reprocessing of the IRAS Survey (IRIS). The SPHERE images in the J , H , or K bands (see Table B.2) are shown in the inset panels with spatial and flux scales optimized for better representation (whereas the actual scales can be appreciated in Fig. 2).

to the entire population of lower-mass stars with a disk (that is, based on [Esplin & Luhman 2019](#), approximately two third of the whole sample of stars with a disk).

2.3. Stellar properties

The 43 sources of this work are divided into three A-, one F-, three G-, nineteen K-, and seventeen M-type stars. To our knowledge, thirteen of the stars are part of multiple systems, with one spectroscopic binary (DQ Tau), five systems with close companions (less than 1'') and seven with further companions. This fraction of multiple stars from our sample (30%) is mildly lower than the overall fraction determined for solar-type stars ($\sim 50\%$, [Raghavan et al. 2010](#)) and significantly lower than that constrained for Taurus ($\sim 70\%$, [Kraus et al. 2011](#)). This is not necessarily a bias since close companions can shorten the disk lifetime or alter their morphology (see [Offner et al. 2023](#), and references therein).

For each target, we retrieved the effective temperature T_{eff} from [López-Valdivia et al. \(2021\)](#) and [Gangi et al. \(2022\)](#), and constructed the spectral energy distribution (SED) through *VizieR*³. We then measured the stellar luminosity L_* by means of a PHOENIX model of the stellar photosphere ([Hauschildt et al. 1999](#)) scaled to the *Gaia* distance and to the R magnitude dereddened by the extinction A_V calculated from the V , R , and I wavebands. Error bars on the stellar luminosity are propagated from d , A_V (20%), and T_{eff} (± 100 K). Finally, we constrained the

stellar mass M_* and age t in the current framework of isochrones through a set of pre-main-sequence tracks ([Siess et al. 2000](#); [Bressan et al. 2012](#); [Baraffe et al. 2015](#); [Choi et al. 2016](#)), including the magnetic tracks by [Feiden \(2016\)](#) that would contrast the tendency of non-magnetic tracks to underestimate stellar mass and age ([Hillenbrand & White 2004](#); [Asensio-Torres et al. 2019](#)). The age interval shown in Table B.1 reflects the variety of values suggested by the different tracks. Some more details on the extraction of the SED and the calculation of the stellar properties are given in Appendix B.

The calculated stellar masses of the individual stellar components span from $2.5 M_{\odot}$ (SU Aur) to $0.4 M_{\odot}$ (8 sources) with 10 intermediate-mass stars ($\geq 1.5 M_{\odot}$), 9 solar-mass stars ($\geq 0.8 M_{\odot}$), and 36 low-mass stars. The age of the majority of the sample is lower than 3 Myr, in line with the known age of the region. Sources that look older from the isochrones are therefore interesting cases that are described in Sect. 4. Here we only point out that the two oldest sources in the sample (CQ Tau and MWC758, 12–14 Myr) are formally not part of Taurus but rather of a nearby association with an estimated age of 18 ± 4 Myr ([Luhman 2023](#)). As for the other sources, the limited statistics and the uncertainty involved prevent us from determining any significant trend between the age of the individual groups by [Luhman \(2023\)](#) and the age that we inferred.

³ <http://vizier.cds.unistra.fr/viz-bin/VizieR>

2.4. Disk properties

From the aforementioned SEDs, we also extracted the IR excess and derived a crude estimate of the dust mass in the disk. For the former, we measured the integrated excess over the stellar photosphere relative to the stellar luminosity in the near-IR (1.2–4.5 μm), mid-IR (4.5–22 μm), and far-IR (22–450 μm), as also shown in Fig. B.1. For the latter, we converted the observed flux at 1.3 mm under standard assumptions (optically thin emission, dust temperature of 20 K, and dust opacity by Beckwith et al. 1990). The derived dust mass should be viewed as a rough estimate due to the simplicity of these assumptions (see e.g., Birnstiel et al. 2018; Zhu 2019; Ribas et al. 2020; Binkert et al. 2021, 2023). The values of excess and disk mass calculated for the entire sample are shown in Table B.2.

Near-, mid-, and far-IR excesses reveal the reprocessing of the stellar photons by the hot, warm, and cold dust at the disk surface or any possible material above it. Empirically, the near-IR excess of a standard, continuous disk is around 10–20% of the stellar luminosity (e.g., Garufi et al. 2022). Lower values point to a disk depleted in the inner regions while higher values to an extra component of hot dust (such as uplifted material in the stellar proximity) reprocessing the stellar light. On the other hand, the far-IR excess typically ranges from 5% for cold, self-shadowed disks to $\gg 50\%$ for disks with heavily exposed outer disk portions. The sample of this work reflects this known variety spanning the near-IR excess from 0% (GM Aur) to 100% (XZ Tau) and the far-IR excess from 0.7% (V1025 Tau) to 57% (DG Tau). Similarly, the dust mass in the disk covers more than two orders of magnitude, from the 1 M_\oplus of V1025 Tau to the 300 M_\oplus of GG Tau.

3. SPHERE observations and data reduction

By construction, all targets in the sample have SPHERE/IRDIS polarimetric images available. All of them have been observed in the H broad-band filter except CQ Tau, LkCa15 (in the J band), MWC758 and CY Tau (in the K). **Observations took place between November 2016 and December 2021. The total integration time spans from 11 to 115 minutes (see Appendix C).** Twelve sources have datasets published by previous authors while the other 31 are unpublished. The complete reference list (publication or observing program) can be found in Tables B.2 and C.1. Among the unpublished datasets, thirteen are from the SPHERE GTO, eleven are from DESTINYs, and seven from various open-time programs.

The data were reduced with the public IRDAP pipeline (IRDIS Data reduction for Accurate Polarimetry, van Holstein et al. 2020) which employs a validated Mueller matrix model to correct for the instrumental polarization and crosstalk of the telescope. Although some of the datasets (e.g., all DESTINYs data) have been acquired in pupil tracking mode to produce a total-intensity image in angular differential imaging (ADI, Marois et al. 2006), in this work we only focus on the polarized intensity. Most recent work used the polarization probed by the radial Stokes parameter Q_ϕ following de Boer et al. (2020):

$$Q_\phi = -Q \cos 2\phi - U \sin 2\phi \quad (1)$$

with ϕ being the azimuthal angle, and Q and U the Stokes parameters from Stokes (1851) that are extracted by IRDAP with the double-difference method (van Holstein et al. 2020). By construction, the Q_ϕ parameter probes the centro-symmetric component of the polarized light, and is equal to the polarized intensity $PI = \sqrt{Q^2 + U^2}$ in the ideal scenario of single scattering from a

face-on disk around a single star. However, several mechanisms can alter this ideal case, such as the presence of companions (Weber et al. 2023), convolution effects from even moderately inclined disks (Heikamp & Keller 2019), or multiple scattering (Canovas et al. 2015). In any of these cases, a fraction of the polarized photons will move from the Q_ϕ parameter to the radial Stokes U_ϕ parameter defined, similarly to the Q_ϕ , as:

$$U_\phi = +Q \sin 2\phi - U \cos 2\phi. \quad (2)$$

In case of strong signal recorded from the U_ϕ parameter, we therefore adopt the PI image instead of the Q_ϕ parameter even though this implies squaring the noise by construction of PI .

The stellar polarization is measured in regions where no bound and background stars nor disks are present, and subsequently corrected. In some datasets, a customized procedure is adopted. This includes the visual selection of the best Stokes parameters from a grid of images with different combinations of stellar polarization, the minimization of negative signal in the Q_ϕ image, and the minimization of positive signal in the U_ϕ image.

The final FITS file produced in our reduction follows a standard format agreed with the Subaru/HiCiao and Gemini Planet Imager communities (see Rich et al. 2022; Benisty et al. 2023) and contains the total intensity I , the Stokes parameters Q and U , and the radial Stokes parameters Q_ϕ and U_ϕ . All images are calibrated in flux by converting the pixel counts to physical units through non-coronagraphic images that are taken in parallel to the science frames. The observed count from these images is equalized to the 2MASS JHK photometry of the relative star, and the conversion factor is applied to the science frames (see Ginski et al. 2022). To obtain a unique value of the disk brightness for each source, we applied the method by Garufi et al. (2014) described in detail by Garufi et al. (2022) and Benisty et al. (2023). The polarized-to-stellar light contrast α_{pol} is a measurement of the fraction of photons released by the star that can reach the disk surface and are effectively scattered toward the observer. It is calculated along the disk major axis to minimize the effect of the scattering phase function and averaged between inner and outer disk radii r_{in} and r_{out} by dividing the radially integrated observed flux $S_{\text{pol}}(r)$ by the stellar flux F_* accounting for its dilution with the separation r :

$$\alpha_{\text{pol}} = \frac{1}{r_{\text{out}} - r_{\text{in}}} \cdot \int_{r_{\text{in}}}^{r_{\text{out}}} S_{\text{pol}}(r) \cdot \frac{4\pi r^2}{F_*} dr. \quad (3)$$

The adoption of the α_{pol} contrast also allows us to minimize the impact of the different filters and integration times t_{int} of the observations in this work. On the one hand, the α_{pol} in the J or K bands (where we have two targets each) is not expected to change from that in the H band unless the dust grains in the disk have a tremendously blue or red color. On the other hand, the α_{pol} is not expected to significantly scale with the integration time while the uncertainty on α_{pol} might have a mild dependence on t_{int} (as is described in Appendix C).

4. Results from the SPHERE images

In this section, we give an overview of the SPHERE images and report on their spatial distribution in Taurus and the correlation with the stellar and disk properties introduced in Sect. 2. A brief description of the individual objects is given in Appendix D.




Fig. 2. Imagery of the sample. The polarization map of all sources is shown with the same angular and flux scales, as indicated in the top-right panel. The Q_ϕ parameter is shown for all sources except GG Tau, XZ Tau, T Tau, and UY Aur for which the PI parameter is shown. All primary stars masked by the coronagraph are indicated by a cyan circle while all secondary stars are indicated by an orange diamond. Each panel is cropped to show all the significant flux detected. The only exception is T Tau, where flux is detected out to the detector edge and for which only the central region is shown. Stars are coarsely sorted by polarized light brightness.

4.1. Overview

The variety of brightness, flux distribution and extent shown by our large sample is appreciable from Fig. 2, where the polarized light from each source is shown in approximate decreasing order of α_{pol} . This contrast spans nearly two orders of magnitude, from $1.6 \cdot 10^{-2}$ (GM Aur) to $3 \cdot 10^{-4}$ (the sources in the small, bottom-right panels of Fig. 2). Four sources (GG Tau, XZ Tau, T Tau, and UY Aur) show very strong signal in the U_ϕ image as well as extended negative signal in the Q_ϕ . These are all systems where the secondary star is in close proximity to the primary (less than $1''$). These companions contribute to the illumination of the circumstellar material, and therefore the observed polarized light is the result of a complex summation of the individual sources of photons (Weber et al. 2023). For these sources, we therefore show the PI image, whereas for all other sources the Q_ϕ image.

Substantially, we can divide the whole sample in three main categories: faint disks, bright disks, and targets where (much of) the scattered light originates from the surrounding environment rather than the disk. The α_{pol} measurement and the category of each source can be found in Table B.2.

4.1.1. Faint disks

By defining a threshold⁴ of $\alpha_{\text{pol}} < 3 \cdot 10^{-3}$ we determine the category of faint and non-detected disks. This is the largest category of the sample (27 sources corresponding to 63% of the sample). Nonetheless, only two of these sources currently have SPHERE publications (CI Tau by Garufi et al. 2020 in the context of a faint disk survey and DH Tau by van Holstein et al. 2021 in a survey of secondary stars with possible polarimetric signal). The class of faint disks can in turn be split between sources where the disk is clearly detected and the basic disk geometry can be inferred (see Appendix D) and those where the polarimetric signal is barely detected. Eleven sources belong to the former sub-category and eleven to the latter. Finally, five disks are formally undetected.

The most notable characteristics of the faint disks is the absence of any obvious substructures from their image. The nearly perfect correspondence between featureless and faint disks indicate that the absence of clear substructures is merely due to our sensitivity and that elusive substructures may still be detectable (see Sect. 5.3). There is however one interesting exception to this trend. MWC480 hosts one of the twelve faintest disks in the sample in terms of contrast but a ring-like structure is clearly visible in the image (see Fig. 2). In fact, in absolute terms the disk is significantly brighter than the others. This is explained by the high stellar luminosity compared to the other faint disks (approximately $20 L_\odot$ against an average $1.2 L_\odot$). It is therefore an illustrative example of the fact that, among disks that scatter few stellar photons, substructures are more easily recovered around bright Herbig stars.

Two other interesting cases are V409 Tau and CW Tau. As is clear from Fig. 2, these stars host an inclined disk. This information was used when determining the photospheric level for the SED (see Appendix B). Both stars are in fact very variable in the visible (around 3 mag based on ASAS-SN curves, Shappee et al. 2014). Based on the disk geometry, we concluded that this variability may largely be due to the disk occultation, and therefore adopted the brightest observed photometry as photospheric level. By doing so, the age calculated from the isochrones turned out to be 3.9 Myr and 3.3 Myr, respectively, which is still older than

the Taurus age (possibly because the stellar flux is always partly extincted) but better than the value of ~ 10 Myr found by adopting an average photometric value. This exercise highlights why the inferred ages of individual systems for which no resolved images are available may differ significantly from the average age of other sources in the host region.

4.1.2. Bright disks

All disks with $\alpha_{\text{pol}} > 3 \times 10^{-3}$ show evidence of substructures. The two old sources of the HD35187 association, MWC758 and CQ Tau, have disks with spiral arms (see Benisty et al. 2015; Hammond et al. 2022). The disk of GG Tau is highly structured with a large cavity, shadows, spirals, and streamers (see Itoh et al. 2014; Keppler et al. 2020). LkCa15 can perhaps be interpreted as a smaller version of GG Tau, exhibiting a thick ring and some inner material within the disk cavity (e.g., Thalmann et al. 2016).

The bowl-shape of the GM Aur disk is clear from the unpublished SPHERE image (see Appendix D), where the disk back side and some spiral-like structures are visible (see also the Subaru image by Oh et al. 2016). Spirals are also detected from the ALMA molecular emission and are probed out to more than 1000 au, suggesting a strong interaction with the cloud (Huang et al. 2021). AB Aur hosts the largest disk of the sample, and possibly of the solar neighborhood. The multiple spiral arms detected from the SPHERE image (Boccaletti et al. 2020) are part of a complex, extended structure that is suggestive of a late-stage infall of material on the star (Tang et al. 2012; Gupta et al. 2023). Similarly, DR Tau shows spiral- and arc-like structures that have been associated with infall from a late encounter with a cloudlet (Mesa et al. 2022). A bright disk with evidence of external infall is also seen in SU Aur (Ginski et al. 2021) where the late accretion process perturbs the disk and is responsible for spirals and shadows. In view of this, the bright disk of GM Aur, AB Aur, DR Tau, and SU Aur could also be classified in the category of ambient material.

4.1.3. Ambient material

Some extended signal not originating from the disk surface is detected from a dozen objects. The prototype of this category is T Tau (e.g., Kasper et al. 2020) where a plethora of extended structures dominates in the polarimetric map and where the disks of the N, Sa, and Sb stars are hardly recognizable. The image of HP Tau shows a similar morphology. In three cases (UX Tau, UY Aur, and XZ Tau), the presence of extended arcs is to be ascribed to the binary interaction. In the prototypical example of UX Tau in particular, a bridge between the A and the C components is detected by SPHERE and by ALMA (Ménard et al. 2020). Some interaction between the individual components is also revealed within the cavity of the bright disk of GG Tau (Keppler et al. 2020).

Signatures of outflow activity or, possibly, disk winds are revealed in four objects. The outflow cavity wall is visible in RY Tau (Garufi et al. 2019; Valegård et al. 2022), DO Tau (Huang et al. 2022), and RW Aur. On top of this, DO Tau also shows very extended arms probing a possible interaction with the nearby HV Tau (Winter et al. 2018). In general, vertically extended material entrained in a disk wind or natal envelope may blend with the disk signal. The net effect is the appearance of a smooth, very flared disk like those observed in DG Tau, RY Tau, and DO Tau (see Fig. 2). These objects are probably revealing the first stages

⁴ This threshold is an approximate median of the whole distribution of α_{pol} calculated (see Garufi et al. 2022).




Fig. 3. Spatial distribution of stellar and disk properties. A simplified version of the map from Fig. 1 is shown to highlight the distribution of bright disks (green), increasingly faint disks (fading to purple), of ambient material (rounded boxes), of binary interaction (rectangular boxes) and of envelope emission (underline), as well of stellar mass (symbol size), and age (symbol color).

of the Class II phase when the optical photons from the star and the disk surface can be observed.

4.2. Statistics and spatial distribution

In Fig. 3, we show the distribution of the various categories across the Taurus region. Evidently, bright and isolated disks in the sample are rare. In fact, stars hosting a disk with substructures and no evidence of interaction with the environment are only 3. Since two of these, CQ Tau and MWC758, are formally not in Taurus, this census is limited to LkCa15. However, if we also include bright disks around binary stars in mutual interaction (two, see Sect. 4.1.3) and around single stars in possible interaction with the environment (four, see Sect. 4.1.2), this count sums up to 7 (16% of the sample). Conversely, targets with faint disks and no sign of interaction are 26 (60%), plus 2 (summing up to 65%) in interaction with the environment. Regardless of the disk brightness, 6 targets (14%) show some interaction with the environment, while 4 sources (9%) show an envelope and detectable outflow activity. Excluding the spectroscopic binary DQ Tau, there are 12 multiple systems in the sample. Of these, 5 (42%) show some interaction among the individual components.

The map of Fig. 3 also reveals a hint of segregation between the central portion of Taurus and the peripheral regions. In the central portion of Taurus where our most represented groups L1495, L1524, and L1536 lies, mostly isolated and faint disks are found. **This segregation cannot be driven by the interstellar extinction since this affects equally the stellar and disk signal considered in the contrast calculation (see Sect. 3). The absolute effect on the observed disk brightness is also minor since in the H band the extinction A_H is only one fifth of the A_V , and the A_V of most of our sample is lower than 2 mag.** The only bright disk in this portion of Taurus that comprises 24 sources is LkCa15 (in the L1536 group at the southern edge of this central region). Interestingly, LkCa15 is also the only star in this main region that appears significantly older (6-9 Myr) from our isochrone calculation. HP Tau and DO Tau are the only sources of the central region in evident interaction with the environment, and both lie on the edges of the region, to the South and to East respectively.

The northern L1517 shows a very different population of objects from our sample, hosting as many as four bright disks out of seven members, with three stars, GM Aur, AB Aur and SU Aur, that are in interaction with the environment (and two of which, AB Aur and SU Aur, in relative proximity). Similarly, the southern portion of Taurus with the L1551, L1558, and T Tau groups have a high incidence of bright disks (with only 2 out of 9 very faint disks), and of ambient material (5 out of 9).

4.3. Correlation with stellar and disk properties

The three categories described in Sect. 4.1 and the disk brightness in scattered light (see Sect. 3) are compared with stellar and disk properties in Fig. 4, and the results are discussed in this section.

4.3.1. Stellar age and mass

Similarly to Fig. 3, any relation between the disk brightness and the stellar properties can be appreciated from Fig. 4 (top panel). From the diagram, it is clear that the majority of stars shows an age in line with Taurus (<3 Myr, e.g. Luhman 2023). On the one hand, most of the young stars are low mass and host a faint disk, with the only bright disk found in GG Tau (a multiple system) and DR Tau (in strong interaction with the environment). On the other hand, a large fraction of the old stars show a bright disk. This dichotomy may suggest that the old age calculated for these sources is real.

Beside MWC758 and CQ Tau (associated with an older group, see Sect. 2), we find 10 sources older than 3 Myr. Two of these, CW Tau and V409 Tau, host an inclined disk that may occult some stellar flux yielding a biased age (see Sect. 4.1). A similar conclusion may in principle be drawn for RW Aur and HP Tau with strong ambient emission. Two of the remaining old stars, DM Tau and V1025 Tau are interesting cases that are discussed below.

It must also be noted that the sample does not contain any young, intermediate-mass star with the notable exception of T Tau. Also, among the few intermediate-mass star we find most (4 out of 5) of the sources in evident interaction with the environment.

4.3.2. Infra-red excess

Garufi et al. (2022) adopted a threshold of 10% to separate high and low IR excesses resulting in a NIR-FIR diagram divided in four quadrants. These quadrants host stars with different disk types. As labeled in Fig. 4 (middle panel), these are transition disks (with both high NIR and low FIR), self-shadowed disks, and evolved disks. Almost all the faint disks of the sample sit in the self-shadowed quadrant. This reflects the geometry where a robust NIR excess (10–20%) originates from the inner disk while small FIR excess (<10%) is reprocessed by the outer disk in penumbra (see Garufi et al. 2020). Conversely, mostly bright disks and sources with ambient emission sit in the upper-right panel.

Nonetheless, a major difference from the large sample by Garufi et al. (2022) arises, that is the very low number of sources with low NIR excess and high FIR excess (top left quadrant). This is by itself an interesting finding as it points to the substantial absence of massive disks with depleted inner regions. The only two objects of the whole quadrant are GM Aur and DM Tau. Instead, to the bottom left we find V1025 Tau with nearly null




Fig. 4. Disk categories across stellar and disk properties. Top: stellar mass vs age. The dashed lines delimit the Taurus age (1–3 Myr, Luhman 2023). Middle: FIR vs NIR excess, shown in fraction of the stellar luminosity. The dashed lines indicate the ideal separation between low and high excesses. Bottom: polarized light contrast vs disk dust mass. The dashed line indicates the detection limit from Garufi et al. (2022).

IR excess (pointing to an evolved Class III target). Interestingly, DM Tau and V1025 Tau are the two old faint disks mentioned in Sect. 4.3.1 suggesting a causality between the two findings that is discussed in Sect. 6.

4.3.3. Disk mass

As is clear from Fig. 4 (bottom panel), there is no tight trend between the disk brightness in scattered light and the dust disk mass. There are many bright and many faint SPHERE images of disks with dust masses higher than $10 M_{\oplus}$. However, disks less massive than $10 M_{\oplus}$ are hardly detected, in agreement with Garufi et al. (2022). In this regard, the detections around GI Tau, CX Tau, HQ Tau, and IP Tau (all with less than $6 M_{\oplus}$) are notable as these are among the least massive disks ever resolved in polarized light. The disk of IP Tau in particular is also relatively bright given the low mass (see also Sect. 5.2).

This diagram also provides a qualitative assessment of why a disk is undetected. On the one hand, disks less massive than $5\text{--}10 M_{\oplus}$ are probably small and would therefore scatter most photons at separations not achievable by the current telescopes. On the other hand, disks more massive than $10 M_{\oplus}$ are most likely extended and their non-detection is due to the self-shadowed geometry of the disk (see Sect. 5.3).

5. Results from the comparison with ALMA

At the time of writing, all the sources of this work except V1025 Tau have been observed with ALMA. More than half of the sample has been observed within the Taurus snapshot survey described by Long et al. (2018, 2019) and Manara et al. (2019). Their sample was built excluding sources with ALMA high-resolution data available at the time. Given that bright, well-known sources were targeted first by the ALMA users, it is not surprising that most of the Long et al. (2019) targets are faint disks in scattered light. In this section, we give a brief review on the continuum ALMA images deferring any comparison with molecular emission to a future work. In particular, we focus on a few interesting cases for which no extensive studies have been reported yet.

5.1. Census of the ALMA sample

Out of the 42 disks with high resolution ALMA data, 9 show a large inner cavity. It is well established that the presence of a large cavity devoid of dust is the primary condition for a strong scattered light signal (see e.g., Garufi et al. 2017) and indeed 7 of these 9 disks with a cavity are bright. These 7 bright disks have been studied in previous ALMA work (Tang et al. 2017; Boehler et al. 2018; Ubeira Gabellini et al. 2019; Facchini et al. 2020; Huang et al. 2020; Francis & van der Marel 2020; Wölfer et al. 2021), and present a geometry where NIR scattered light is detected within the ALMA cavity, revealing spirals (in CQ Tau, MWC758, GG Tau, AB Aur, and GM Aur, Hammond et al. 2022; Benisty et al. 2015; Keppler et al. 2020; Boccaletti et al. 2020, and Appendix D, respectively) or some degree of asymmetry (Thalmann et al. 2016; Ménard et al. 2020, in LkCa15 and UX Tau). However, the two faint disks with an ALMA cavity (IP Tau and DM Tau) are also interesting cases that are described in Sect. 5.2.

Out of the remaining 33 disks with no large cavity, 9 disks show obvious evidence of substructures in the currently available ALMA data. Seven of these 9 are faint disks that are described




Fig. 5. Disks with an ALMA cavity that are faint in scattered light. The ALMA image (in red) of IP Tau and DM Tau is compared to the high-pass filtered SPHERE image (in blue). The innermost region of the original SPHERE images is masked by the coronagraph.

in Sect. 5.3. The remaining two (DG Tau and RY Tau) are the prototypical example of partly embedded disks (see Sect. 4.1.3) for which the NIR images only yields limited information on the disk surface. They are showcases of the early formation of substructures in disks (as in the pioneering example of HL Tau, ALMA Partnership et al. 2015).

As many as 24 ALMA disks can be coarsely considered featureless at the time being, although the diverse angular resolution and sensitivity of the various datasets nullify any quantitative assessment of the class of objects. Future improved ALMA observations or reanalysis of the current sample (see e.g., Jennings et al. 2022; Zhang et al. 2023) may reveal further substructures. All these disks are either faint in scattered light or dominated by ambient emission. Eight of these disks are part of a multiple system as described in Sect. 5.4. Some additional information about the individual cases are given in Appendix D.

5.2. The peculiar cases of IP Tau and DM Tau

The two disks with a cavity from the ALMA continuum that appear faint from our SPHERE observations are IP Tau and DM Tau. They are both per se interesting cases, as described in Sect. 4.3. As is clear from Fig. 4, DM Tau is the only low-mass star that looks older than 3 Myr⁵, and is among the few Class II sources ever observed with null NIR excess that is not particularly bright in scattered light. IP Tau is the brightest disk in the sample with less than 10 M_{\oplus} in dust mass (see Fig. 4), and is probably among the brightest ever observed of such a type.

In Fig. 5, we compare the SPHERE image of IP Tau and DM Tau with the ALMA continuum by Long et al. (2019) and Hashimoto et al. (2021). We show a high-pass filtered version of the SPHERE image obtained by convolving the image with a gaussian profile, and then removing the convolved image from the original image. This is a common technique to highlight elusive substructures. This exercise clearly reveals the presence of spirals in the disk of IP Tau that are co-spatial with the inner edge of the ALMA disk. In DM Tau, a clear asymmetry is visible to the SE where the inner edge of the NIR emission seems to penetrate deeper in the ALMA cavity with respect to the symmetrical portion around the minor axis, to NW. In principle, this asymmetry may also point to an unresolved feature like a spiral. These findings are in line with the recurrent presence of such

substructures in disks with a cavity described above (see also van der Marel et al. 2021), and are further discussed in Sect. 6.

5.3. The cases of the self-shadowed disks

In Fig. 6, we compare ALMA and SPHERE images of the seven faint disks in our sample that are classified as substructured disks by Long et al. (2019). Generally speaking, the comparison reveals that only a fraction of the disk is probed by the SPHERE observations. This is particularly true for CI Tau (see also Garufi et al. 2022) and DN Tau. This configuration is the most direct proof that these disks are self-shadowed, like formulated by Dullemond & Dominik (2004) and shown observationally by Garufi et al. (2022).

Other examples in Fig. 6 such as UZ Tau and IQ Tau are similar. The SPHERE signal extent is comparable to ALMA but it is reasonable to assume that the gas disk extent is larger than the ALMA continuum and is therefore also not probed by our observations. In the case of DL Tau, as in MWC480, some ring-like structures do become visible after light smoothing of the image and aided by the comparison with ALMA. Finally, in the case of DS Tau the comparison with ALMA allows to determine the presence of a weak Q_{ϕ} signal consistent with disk scattered light, even though this target is formally treated as a non-detection in this work (see Appendix D).

5.4. The cases of multiple systems

Only three of our disks in multiple systems are more extended than 0.2'' from the ALMA continuum (UX Tau A, UZ Tau, and V710 Tau). In each of these three systems, the separation between the individual components is large ($>2.7''$). These three disks are clearly resolved with SPHERE. Furthermore, a disk is also resolved around UX Tau C (see Ménard et al. 2020) and UZ Tau B and C (see Fig. 2), where two disks with the same position angle of the primary disk (in line with the ALMA image by Manara et al. 2019) are resolved out to 0.4'' and 0.2'', respectively.

In the remaining eight cases, ALMA constrains a small disk ($<0.15''$) with a steep decay at the outer radius that is indicative of tidal truncation (Manara et al. 2019). In only one of these cases, DH Tau A, the disk is clearly resolved by SPHERE (beside the detection of the disk around DH Tau B by van Holstein et al. 2021) while in one case, RW Aur, some disk emission is embedded with some ambient material (see Sect. 4.1.3). In all other cases the disk is not detected because either hidden by the interacting material (XZ Tau, UY Aur, T Tau) or because no significant polarimetric signal is visible (HN Tau, V807 Tau, DK Tau).

6. Discussion

This work contributes to the general effort of the planet-forming disk community to characterize less and less exceptional objects and to study the disk demography as a whole. Thanks to the work put in place by many authors, a very large fraction of the observable Taurus population has now been observed. In this work, we show that the current sample is relatively complete for stars with mass above 0.4 M_{\odot} . It does however also urge the community to develop near-IR facilities that can probe optically fainter stars as this means covering more embedded, lower-mass, and further objects.

One of the key-results of this work is that the prototypical disk in Taurus (1–3 Myr) appears faint in scattered light. This

⁵ excluding V409 Tau for which the disk inclination may bias the age estimate (see Sect. 2.3).




Fig. 6. Self-shadowed disks in the sample. The ALMA image of those disks that show evidence of substructures is placed beside an intentionally saturated SPHERE image obtained with a light smoothing. The ALMA beam is shown to the bottom right of each image, the physical scale (0.5'') common to ALMA and SPHERE relative panels to the bottom left. All ALMA images are from Long et al. (2019) except CI Tau from Rosotti et al. (2021).

is partly due to the small physical extent of some disks and partly to the low incidence of an inner cavity in these disks that would allow the efficient illumination of the outer disk regions. In fact, the partition between disks with and without a large cavity from ALMA images naturally demarcates our boundary between bright and faint disks from the SPHERE images. A large fraction of these faint disks will never form an inner cavity, as suggested by complete Spitzer survey and ALMA demographical studies (Williams & Cieza 2011; Cieza et al. 2019, 2021, respectively). Nonetheless, the fraction of bright disks from surveys of Class II sources only is still expected to increase with time because disks that do not form cavity will turn into evolved Class III sources, leaving only disks with the cavity in the sample.

In this work, we also find a possible spatial segregation for different disk types within the whole region, with the central portion of Taurus mainly hosting faint, isolated disks and with the Taurus peripheries hosting a higher number of interacting, bright disks. It is tempting to draw an evolutionary history for the region that is however not captured yet by the estimated ages of the individual Taurus groups. It could instead be linked to the higher stellar mass of some peripheral group members (see e.g., SU Aur, AB Aur or T Tau and UX Tau) for which a higher incidence of interaction with the environment is found in this work or a faster disk evolution is reasonably expected.

Understanding the disk demography from a uniform sample like this in Taurus also benefits by studying the outliers and the extreme cases. A visual synopsis of the peculiar objects encountered in this work is given in Fig. 7. The main box is reserved to the large portion of faint disks in the sample. The disk extent constrained from the ALMA continuum or the dust mass measured from the millimeter photometry aid the interpretation of the actual disk geometry. On the one hand extended, self-shadowed disks are well represented in the Taurus sample with the most extreme cases being MWC480 and DL Tau (that are very massive but are relatively easily detected by SPHERE) or DN Tau and DQ Tau (that are barely detected by SPHERE but are only mildly massive). On the other hand, small disks that are less than $10 M_{\oplus}$ in dust mass are hardly detectable by SPHERE. The resolved detection of as many as four disks less massive than that (HQ, CX, GI, and IP Tau) is an important step ahead toward the characterization of less exceptional objects.

On the opposite end of the dust mass distribution we find the well-known giant disks such as that of GG Tau and GM Aur. This work highlights their low incidence since the sample bias is alleviated, and shows that their fraction is no larger than 20%. This fraction reduces to just 5% if we only consider isolated, bright disks and exclude the two old members of a nearby association MWC758 and CQ Tau. It is tempting and possibly reasonable to consider these disks the only survivors of an ancient disk population with an estimated age of 18 Myr. In any case, the fraction of bright disks could be even smaller when the population of lower-mass stars not observed by SPHERE is considered.

It is increasingly clear to the community that planet-forming disks are not isolated structures. The incidence of targets that are interacting with the environment is another pivotal element in disk evolution studies that can only be constrained by looking at an unbiased sample. The 30% fraction that is measured in this work could however be smaller when lower-mass stars are considered. Indeed, we find that the majority of stars interacting with their environment are of intermediate mass, although this may be partly due to the higher luminosity of these stars that can reveal ambient material on a larger volume. In addition, a




Fig. 7. Visual synopsis of the sample. The demography of the SPHERE sample studied in this work is summarized with illustrative examples. The main central box includes the prototypical object of the sample, that is a faint, isolated disk. All the peculiar cases that do not belong to this category are depicted in the smaller boxes. The name of the source shown is underlined.

large fraction (40%) of stars in multiple systems show mutual interactions.

Finally, it is illustrative to look at the beginning and end stages of evolution within our sample of Class II. The morphology observed in objects like DG Tau and RY Tau suggests that we are seeing young Class II objects where optical photons from the star and the disk can reach the observer, although still blended with the contribution from natal envelope and outflows. At the other extreme, we propose that two objects of the sample, DM Tau and IP Tau, represent the final stage of the Class II phase. Both are relatively faint in scattered light despite the presence of a large ALMA cavity. DM Tau shows no NIR excess from the SED despite an inner disk component that is detected by ALMA (Hashimoto et al. 2021), pointing to a population of large grains only in the inner region. It is also the only low-mass star in the sample that appears to be older than 3 Myr. Given the small uncertainties relative to this measurement (no extinction, precise distance), we consider this estimate correct. IP Tau hosts a disk with a very small amount of dust but, in this regard, it is still relatively bright in scattered light. The presence of spirals detected in the SPHERE image reveals complex dynamical processes as in more massive, brighter transition disks (van der Marel et al. 2021), but the low disk mass compared to the extent of the scattered light emission suggests that this object is transitioning to the Class III stage, like the only such a case in our sample of V1025 Tau.

7. Summary

In this work, we examine the demographics of planet-forming disks in Taurus through the full SPHERE dataset available. This sums up to 43 sources with near-IR polarimetric images (31 un-

published) that probe the scattered light from any circumstellar material. Our main findings are the following.

- The available sample covers approximately one fifth of the total Class II population in Taurus, and approximately a half of those that can be observed with SPHERE (three quarters of those observable in optimal conditions). All in all, the sample is quite complete above $0.4 M_{\odot}$ but is completely blind to lower-mass stars.
- Thirteen of the 43 targets are multiple systems. The individual components in the sample divide in 10 intermediate-mass, 9 solar-mass, and 36 low-mass stars.
- The majority of the sample (63%) are faint disks with no prior study from the literature. No obvious substructures are visible in these disks at the current sensitivity and resolution limits. In a few cases, rings that are clear from ALMA are recovered in the SPHERE images (e.g., MWC480, DL Tau).
- Disks that are less massive than $10 M_{\oplus}$ are rarely detected by SPHERE because of their limited size. However, several massive disks are also barely detectable (e.g., DN Tau). These are the extended self-shadowed disks that have not (yet) formed a cavity. Their high occurrence in Taurus is likely due to their young age.
- Ambient material is visible in 30% of the sample, with a mild prevalence in intermediate-mass stars. This class includes disks in interaction with the environment (e.g., SU Aur), binaries in mutual interaction (e.g., UY Aur), and stars with evidence of natal envelope and outflow activities (e.g., RY Tau).
- About 40% of the binary systems show some evidence of interaction between the individual components. The disks are rarely detected in these systems (and never in case of close companions). This can be reconciled with small disk sizes

due to truncation proposed from previous ALMA observations.

- Bright, isolated disks like e.g., LkCa15 are rare (5%). Bright disks in general (including those in interaction with environment and companions) sum up to 23%. Only two of these disks, GM Aur and DM Tau, are significantly depleted of near-IR excess from the SED. This incidence is lower than in previous study with diversified samples and points to an early evolutionary stage of Taurus sources.
- The presence of a disk cavity from ALMA naturally intercepts all and only bright disks in the near-IR. Two interesting exceptions are the relatively faint disks of IP Tau and DM Tau, showing mild spirals and asymmetries like the other disks with an ALMA cavity. The low dust mass of the former and the null near-IR excess and old age of the latter suggest that these disks are transitioning to an evolved stage.
- A hint of spatial segregation for disk types is visible within the Taurus region. The central portion of the region primarily hosts faint, isolated disks while a higher occurrence of bright, interacting disks are found in the Taurus periphery.
- Some stars with an inclined disk appear artificially older (e.g., V409 Tau) because of occultation of the stellar photosphere. Apart from these, it is clear that brighter disks are found around older stars. This indicates that the ages derived from isochrones are probably reasonably accurate and point to an expected evolutionary path where mostly only disks with a large cavity survive more than 3 Myr.

In summary, several findings indicate an early evolutionary stage for the overall Taurus population, although some individual objects may represent advanced or exceptional evolutionary stages and some others may reveal intermediate phases of the disk lifetime. Together with the parallel work on Chamaeleon (Ginski et al.) and Orion (Valegard et al.), this is a first effort to characterize the demographics of planet-forming disks of an individual region from near-IR images. Future work includes the characterization of additional star-forming regions as well as the direct comparison of the results found in different regions to determine the evolution of planet-forming disks from the largest sample currently possible.

Acknowledgements. We thank the referee for the useful comments as well as F. Long, G. Rosotti, J. Hashimoto, and N. van der Marel for sharing their ALMA data. SPHERE is an instrument designed and built by a consortium consisting of IPAG (Grenoble, France), MPIA (Heidelberg, Germany), LAM (Marseille, France), LESIA (Paris, France), Laboratoire Lagrange (Nice, France), INAF Osservatorio di Padova (Italy), Observatoire de Genève (Switzerland), ETH Zurich (Switzerland), NOVA (Netherlands), ONERA (France) and ASTRON (Netherlands) in collaboration with ESO. SPHERE was funded by ESO, with additional contributions from CNRS (France), MPIA (Germany), INAF (Italy), FINES (Switzerland) and NOVA (Netherlands). SPHERE also received funding from the European Commission Sixth and Seventh Framework Programmes as part of the Optical Infrared Coordination Network for Astronomy (OPTICON) under grant number RII3-Ct-2004-001566 for FP6 (2004–2008), grant number 226604 for FP7 (2009–2012) and grant number 312430 for FP7 (2013–2016). This research has made use of the VizieR catalogue access tool, CDS, Strasbourg, France (DOI: 10.26093/cds/vizieR). The original description of the VizieR service was published in 2000, A&AS 143, 23. This work was supported by the PRIN-INAF 2019 Planetary Systems At Early Ages (PLATEA). We also acknowledge financial support from the Programme National de Planétologie (PNP) and the Programme National de Physique Stellaire (PNPS) of CNRS-INSU in France and from ANID – Millennium Science Initiative Program – Center Code NCN2021080. This project has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (PROTOPLANETS, grant agreement No. 101002188). C.F.M. is funded by the European Union (ERC, WANDA, 101039452). S.P. acknowledges support from FONDECYT grant 1231663. P.W. acknowledges support from FONDECYT grant 3220399. S.F. is funded by the European Union under the European Union’s Horizon Europe Research & Innovation Programme 101076613 (UNVEIL). Views and opinions expressed are however those of the

author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them. This project has received funding from the European Research Council (ERC) under the European Union’s Horizon Europe research and innovation programme (grant agreement No. 101053020, project Dust2Planets).

References

- Akeson, R. L., Jensen, E. L. N., Carpenter, J., et al. 2019, *ApJ*, 872, 158
 Alcalá, J. M., Gangi, M., Biazzo, K., et al. 2021, *A&A*, 652, A72
 ALMA Partnership, Brogan, C. L., Pérez, L. M., et al. 2015, *ApJ*, 808, L3
 Andrews, S. M. 2020, *ARA&A*, 58, 483
 Ansdell, M., Williams, J. P., van der Marel, N., et al. 2016, *ApJ*, 828, 46
 Asensio-Torres, R., Currie, T., Janson, M., et al. 2019, *A&A*, 622, A42
 Avenhaus, H., Quanz, S. P., Garufi, A., et al. 2018, *The Astrophysical Journal*, 863, 44
 Baraffe, I., Homeier, D., Allard, F., & Chabrier, G. 2015, *A&A*, 577, A42
 Beckwith, S. V. W., Sargent, A. I., Chini, R. S., & Guesten, R. 1990, *AJ*, 99, 924
 Benisty, M., Dominik, C., Follette, K., et al. 2023, in *Astronomical Society of the Pacific Conference Series*, Vol. 534, *Astronomical Society of the Pacific Conference Series*, ed. S. Inutsuka, Y. Aikawa, T. Muto, K. Tomida, & M. Tamura, 605
 Benisty, M., Juhasz, A., Boccaletti, A., et al. 2015, *A&A*, 578, L6
 Beuzit, J. L., Vigan, A., Mouillet, D., et al. 2019, *A&A*, 631, A155
 Binkert, F., Szulágyi, J., & Birnstiel, T. 2021, *MNRAS*, 506, 5969
 Binkert, F., Szulágyi, J., & Birnstiel, T. 2023, *MNRAS*, 523, 55
 Birnstiel, T., Dullemond, C. P., Zhu, Z., et al. 2018, *ApJ*, 869, L45
 Boccaletti, A., Di Folco, E., Pantin, E., et al. 2020, *A&A*, 637, L5
 Boehler, Y., Ricci, L., Weaver, E., et al. 2018, *ApJ*, 853, 162
 Bressan, A., Marigo, P., Girardi, L., et al. 2012, *MNRAS*, 427, 127
 Canovas, H., Ménard, F., de Boer, J., et al. 2015, *A&A*, 582, L7
 Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
 Choi, J., Dotter, A., Conroy, C., et al. 2016, *ApJ*, 823, 102
 Cieza, L. A., González-Ruilova, C., Hales, A. S., et al. 2021, *MNRAS*, 501, 2934
 Cieza, L. A., Ruiz-Rodríguez, D., Hales, A., et al. 2019, *MNRAS*, 482, 698
 de Boer, J., Langlois, M., van Holstein, R. G., et al. 2020, *A&A*, 633, A63
 Dohlen, K., Langlois, M., Saisse, M., et al. 2008, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 7014, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 3
 Dullemond, C. P. & Dominik, C. 2004, *A&A*, 417, 159
 Esplin, T. L. & Luhman, K. L. 2019, *AJ*, 158, 54
 Facchini, S., Benisty, M., Bae, J., et al. 2020, *A&A*, 639, A121
 Facchini, S., van Dishoeck, E. F., Manara, C. F., et al. 2019, *arXiv e-prints*, arXiv:1905.09204
 Feiden, G. A. 2016, *A&A*, 593, A99
 Fitton, S., Tofflemire, B. M., & Kraus, A. L. 2022, *Research Notes of the American Astronomical Society*, 6, 18
 Francis, L. & van der Marel, N. 2020, *ApJ*, 892, 111
 Gaia Collaboration. 2016, *VizieR Online Data Catalog*, 1337
 Gaia Collaboration, Vallenari, A., Brown, A. G. A., et al. 2022, *arXiv e-prints*, arXiv:2208.00211
 Galli, P. A. B., Loinard, L., Bouy, H., et al. 2019, *A&A*, 630, A137
 Galli, P. A. B., Loinard, L., Ortiz-Léon, G. N., et al. 2018, *ApJ*, 859, 33
 Gangi, M., Antonucci, S., Biazzo, K., et al. 2022, *A&A*, 667, A124
 Garufi, A., Benisty, M., Pinilla, P., et al. 2018, *A&A*, 620, A94
 Garufi, A., Dominik, C., Ginski, C., et al. 2022, *A&A*, 658, A137
 Garufi, A., Meeus, G., Benisty, M., et al. 2017, *A&A*, 603, A21
 Garufi, A., Podio, L., Bacciotti, F., et al. 2019, *A&A*, 628, A68
 Garufi, A., Podio, L., Codella, C., et al. 2020, *A&A*, 636, A65
 Garufi, A., Quanz, S. P., Schmid, H. M., et al. 2014, *A&A*, 568, A40
 Garufi, A., Quanz, S. P., Schmid, H. M., et al. 2016, *A&A*, 588, A8
 Ginski, C., Facchini, S., Huang, J., et al. 2021, *ApJ*, 908, L25
 Ginski, C., Gratton, R., Bohn, A., et al. 2022, *A&A*, 662, A74
 Ginski, C., Stolker, T., Pinilla, P., et al. 2016, *A&A*, 595, A112
 Gupta, A., Miotello, A., Manara, C. F., et al. 2023, *A&A*, 670, L8
 Hammond, I., Christiaens, V., Price, D. J., et al. 2022, *MNRAS*, 515, 6109
 Hashimoto, J., Dong, R., Kudo, T., et al. 2012, *ApJ*, 758, L19
 Hashimoto, J., Muto, T., Dong, R., et al. 2021, *ApJ*, 911, 5
 Hauschildt, P. H., Allard, F., & Baron, E. 1999, *ApJ*, 512, 377
 Heikamp, S. & Keller, C. U. 2019, *A&A*, 627, A156
 Hillenbrand, L. A. & White, R. J. 2004, *ApJ*, 604, 741
 Huang, J., Andrews, S. M., Dullemond, C. P., et al. 2020, *ApJ*, 891, 48
 Huang, J., Bergin, E. A., Öberg, K. I., et al. 2021, *ApJS*, 257, 19
 Huang, J., Ginski, C., Benisty, M., et al. 2022, *ApJ*, 930, 171
 Itoh, Y., Oasa, Y., Kudo, T., et al. 2014, *Research in Astronomy and Astrophysics*, 14, 1438

- Jennings, J., Booth, R. A., Tazzari, M., Clarke, C. J., & Rosotti, G. P. 2022, *MNRAS*, 509, 2780
- Joncour, I., Duchêne, G., Moraux, E., & Motte, F. 2018, *A&A*, 620, A27
- Kasper, M., Santhakumari, K. K. R., Herbst, T. M., et al. 2020, *A&A*, 644, A114
- Kenyon, S. J., Gómez, M., & Whitney, B. A. 2008, in *Handbook of Star Forming Regions*, Volume I, ed. B. Reipurth, Vol. 4, 405
- Keppler, M., Penzlin, A., Benisty, M., et al. 2020, *A&A*, 639, A62
- Köhler, R., Kasper, M., Herbst, T. M., Ratzka, T., & Bertrang, G. H. M. 2016, *A&A*, 587, A35
- Kraus, A. L., Ireland, M. J., Martinache, F., & Hillenbrand, L. A. 2011, *ApJ*, 731, 8
- Krolikowski, D. M., Kraus, A. L., & Rizzuto, A. C. 2021, *AJ*, 162, 110
- Lada, C. J. 1987, in *IAU Symposium*, Vol. 115, *Star Forming Regions*, ed. M. Peimbert & J. Jugaku, 1–17
- Laws, A. S. E., Harries, T. J., Setterholm, B. R., et al. 2020, *ApJ*, 888, 7
- Leike, R. H., Glatzle, M., & Enßlin, T. A. 2020, *A&A*, 639, A138
- Long, F., Herczeg, G. J., Harsono, D., et al. 2019, *ApJ*, 882, 49
- Long, F., Pinilla, P., Herczeg, G. J., et al. 2018, *ApJ*, 869, 17
- López-Valdivia, R., Sokal, K. R., Mace, G. N., et al. 2021, *ApJ*, 921, 53
- Luhman, K. L. 2023, *AJ*, 165, 37
- Macintosh, B., Graham, J. R., Ingraham, P., et al. 2014, *Proceedings of the National Academy of Science*, 111, 12661
- Manara, C. F., Tazzari, M., Long, F., et al. 2019, *A&A*, 628, A95
- Marois, C., Lafrenière, D., Doyon, R., Macintosh, B., & Nadeau, D. 2006, *ApJ*, 641, 556
- Mayama, S., Hashimoto, J., Muto, T., et al. 2012, *ApJ*, 760, L26
- Ménard, F., Cuello, N., Ginski, C., et al. 2020, *A&A*, 639, L1
- Mesa, D., Ginski, C., Gratton, R., et al. 2022, *A&A*, 658, A63
- Offner, S. S. R., Moe, M., Kratter, K. M., et al. 2023, in *Astronomical Society of the Pacific Conference Series*, Vol. 534, *Astronomical Society of the Pacific Conference Series*, ed. S. Inutsuka, Y. Aikawa, T. Muto, K. Tomida, & M. Tamura, 275
- Oh, D., Hashimoto, J., Carson, J. C., et al. 2016, *ApJ*, 831, L7
- Pascucci, I., Testi, L., Herczeg, G. J., et al. 2016, *ApJ*, 831, 125
- Price, D. J. 2007, *PASA*, 24, 159
- Raghavan, D., McAlister, H. A., Henry, T. J., et al. 2010, *ApJS*, 190, 1
- Ribas, Á., Espaillat, C. C., Macías, E., & Sarro, L. M. 2020, *A&A*, 642, A171
- Rich, E. A., Monnier, J. D., Aarnio, A., et al. 2022, *AJ*, 164, 109
- Roccatagliata, V., Franciosini, E., Sacco, G. G., Randich, S., & Sicilia-Aguilar, A. 2020, *A&A*, 638, A85
- Rosotti, G. P., Ilee, J. D., Facchini, S., et al. 2021, *MNRAS*, 501, 3427
- Shappee, B. J., Prieto, J. L., Grupe, D., et al. 2014, *ApJ*, 788, 48
- Siess, L., Dufour, E., & Forestini, M. 2000, *A&A*, 358, 593
- Soler, J. D., Zucker, C., Peek, J. E. G., et al. 2023, *arXiv e-prints*, arXiv:2305.13180
- Stokes, G. G. 1851, *Transactions of the Cambridge Philosophical Society*, 9, 399
- Tang, Y.-W., Guilloteau, S., Dutrey, A., et al. 2017, *ApJ*, 840, 32
- Tang, Y. W., Guilloteau, S., Piétu, V., et al. 2012, *A&A*, 547, A84
- Thalmann, C., Janson, M., Garufi, A., et al. 2016, *ApJ*, 828, L17
- Ubeira Gabellini, M. G., Miotello, A., Facchini, S., et al. 2019, *MNRAS*, 486, 4638
- Ueda, T., Kataoka, A., & Tsukagoshi, T. 2022, *ApJ*, 930, 56
- Valegård, P. G., Ginski, C., Dominik, C., et al. 2022, *A&A*, 668, A25
- van der Marel, N., Birmstiel, T., Garufi, A., et al. 2021, *AJ*, 161, 33
- van Holstein, R. G., Girard, J. H., de Boer, J., et al. 2020, *A&A*, 633, A64
- van Holstein, R. G., Stolker, T., Jensen-Clem, R., et al. 2021, *A&A*, 647, A21
- Weber, P., Pérez, S., Guidi, G., et al. 2023, *MNRAS*, 518, 5620
- Williams, J. P. & Cieza, L. A. 2011, *ARA&A*, 49, 67
- Winter, A. J., Booth, R. A., & Clarke, C. J. 2018, *MNRAS*, 479, 5522
- Wölfer, L., Facchini, S., Kurtovic, N. T., et al. 2021, *A&A*, 648, A19
- Zhang, S., Kalscheur, M., Long, F., et al. 2023, *arXiv e-prints*, arXiv:2305.03862
- Zhu, Z. 2019, *MNRAS*, 483, 4221
- ⁸ Departamento de Física, Universidad de Santiago de Chile, Avenida Victor Jara 3659, Santiago, Chile
- ⁹ Mullard Space Science Laboratory, University College London, Holmbury St Mary, Dorking, Surrey RH5 6NT, UK
- ¹⁰ Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge, CB3 0HA, UK
- ¹¹ Institute for Astronomy, University of Hawaii, Hawaii, USA
- ¹² Instituto de Estudios Astrofísicos, Facultad de Ingeniería y Ciencias, Universidad Diego Portales, Avenida Ejército Libertador 441, Santiago, Chile
- ¹³ Anton Pannekoek Institute for Astronomy, University of Amsterdam, Amsterdam, The Netherlands
- ¹⁴ Dipartimento di Fisica, Università degli Studi di Milano, Via Celoria 16, 20133 Milano, Italy
- ¹⁵ Department of Astronomy, Columbia University, 538 W. 120th Street, Pupin Hall, New York, NY, United States of America
- ¹⁶ Department of Astronomy, University of Florida, Gainesville, FL 32611, USA
- ¹⁷ Observatoire de Genève, Université de Genève, 51 Ch. des Maillettes, 1290 Sauverny, Switzerland
- ¹⁸ Max Planck Institute for Astronomy, Königstuhl 17, D-69117 Heidelberg, Germany
- ¹⁹ Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands
- ²⁰ Department of Astronomy, Stockholm University, 106 91 Stockholm, Sweden
- ²¹ INAF - Osservatorio Astrofisico di Catania, Via S.Sofia 78, I-95123 Catania, Italy
- ²² Department of Astronomy, The University of Michigan West Hall 323, 1085 S. University Avenue Ann Arbor, MI 48109
- ²³ School of Physics and Astronomy, Monash University, Vic 3800, Australia
- ²⁴ ETH Zürich, Department of Physics, Wolfgang-Pauli-Strasse 27, CH-8093, Zürich, Switzerland
- ²⁵ Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, 35122 Padova, Italy

¹ INAF, Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, I-50125 Firenze, Italy
e-mail: antonio.garufi@inaf.it

² Centre for Astronomy, Dept. of Physics, National University of Ireland Galway, University Road, Galway H91 TK33, Ireland

³ European Southern Observatory, Alonso de Córdova 3107, Casilla 19001, Vitacura, Santiago, Chile

⁴ Laboratoire Lagrange, Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, Nice Cedex 4, France

⁵ Univ. Grenoble Alpes, CNRS, IPAG, 38000 Grenoble, France

⁶ European Southern Observatory (ESO), Garching bei München, Germany

⁷ Millennium Nucleus on Young Exoplanets and their Moons (YEMS)

Appendix A: Spatial distribution of the sample

The spatial distribution of the entire Taurus sample by Luhman (2023) is shown in Fig. A.1. The sample of this work is also shown in the map together with the Class II sources that are possibly observable by SPHERE. The names of the groups with at least one member in this work are also shown.




Fig. A.1. The sample of this work compared with the whole Taurus. The spatial distribution of all Taurus sources from Luhman (2023) is compared with the that of Class II sources, of Class II sources that are observable by SPHERE, and of those observed. The groups with at least one member in this work are labeled (see Luhman 2023, for the complete list)

Appendix B: SEDs and properties of the sample

The SEDs of the entire sample described in Sect. 2.3 are shown in Fig. B.1. In the following, we describe a few individual cases that requires an in-depth analysis. The stellar and disk properties described in Sects. 2.3 and 2.4 are listed in Tables B.1 and B.2.

HN Tau. There is no value of extinction A_V , scaled with the Cardelli et al. (1989) law, that fit the entire optical photometry (0.3 to 0.9 μm). To avoid introducing the high uncertainty on the deriving properties, we do not include this source in the statistical analysis. In fact, even scaling the model to the near-IR photometry (less affected by the extinction) the resulting age calculated from isochrones turned out to be of the order of 10 Myr. In line with our finding, Alcalá et al. (2021) defined this object as subluminoous explaining it with obscuration from the inclined disk (Long et al. 2019). We also note that this source is one of the eight with a very poor *Gaia* solution (see Sect. 2.3), indicating clear problems in determining the photocenter.

V409 Tau and CW Tau. As commented in Sect. 4.1, these stars are very variable in the visible. The origin of the variability is necessary to determine what level of observed photometry to adopt as photospheric level. Here, we took advantage of the notion that their disk is very inclined (see Fig. 2 as well as Long et al. 2019 and Ueda et al. 2022) to conclude that the variability is due to partial disk occultation and therefore adopt the brightest observed photometry as photospheric level. By doing so, the near-IR photometry lies slightly below the stellar model but this is probably due to the extinction still significant at these wavelengths.

Appendix C: Observing setup

The setup of all the observations in this work is given in Table C.1. The individual detector integration time (DIT) spans from the 0.837 sec of the non-coronagraphic observations of V807 Tau to the 96 sec of DM Tau. However, most of the targets have been observed with a DIT of 32 sec or 64 sec.

Instead, a larger variety of total integration times t_{int} is present spanning this from 11 minutes for several sources from the GTO to the >100 minutes of V710 Tau and UZ Tau. The adoption of the contrast α_{pol} to assess the disk brightness allows us to minimize this diversity. In fact, the uncertainty on this value (tabulated in Table B.2) only mildly depends on the t_{int} . Sources with $\Delta\alpha_{\text{pol}} \lesssim 3 \cdot 10^{-4}$ have been observed on average for 46 minutes, those with $\Delta\alpha_{\text{pol}} = 3 \cdot 10^{-4}$ for 32 minutes, and those with $\Delta\alpha_{\text{pol}} \gtrsim 3 \cdot 10^{-4}$ for 21 minutes. Since the typical detection limit of a disk is $\alpha_{\text{pol}} = 3 \cdot 10^{-4}$ (see also Garufi et al. 2022), we conclude that the diverse integration times of the observations in this work do not have a significant impact on demographic results of this work. An illustrative example of this is that of CY Tau ($t_{\text{int}} = 51$ min) and CX Tau ($t_{\text{int}} = 11$ min). The former has a much better $\Delta\alpha_{\text{pol}}$ (10^{-4} vs $3 \cdot 10^{-4}$) but the α_{pol} is comparable ($6 \cdot 10^{-4}$ vs $4 \cdot 10^{-4}$) meaning that the disk of CY Tau is observed with a better signal-to-noise than that of CX Tau but it is assessed to be equally faint.

Appendix D: Notes on SPHERE images with no prior publications

Here we briefly describe the SPHERE images with no prior publications available. Sources that are not mentioned are published like referenced in Table B.2. The basic constraints on the disk geometry extractable from the SPHERE images (outermost separation with detectable signal, crude inclination, and position angle) are listed in Table D.1. In the following, we refer to a QU pattern as the expected alignment of positive and negative signal in the Q and U image in case of centro-symmetric scattering, and that can be used to infer a disk detection where the signal detected in the Q_ϕ image is of dubious origin (see e.g., Appendix C in Garufi et al. 2022). Furthermore, the disk extent constrained by ALMA is from the continuum emission unless otherwise specified.

RW Aur. The Q_ϕ image reveals a peculiar morphology around the primary star. The QU pattern indicates the detection of a disk with position angle of approximately 45° . However, bright signal extended in the direction orthogonal to the disk is also detected, and this likely probes the two outflow cavities in a morphology similar to RY Tau (Garufi et al. 2019). Interestingly, a narrow dark lane resembling a gap sculpted by a collimated jet is visible to the SE. Since the disk signal is blended with the outflow material (see Sect. 4.1.3), the calculation of α_{pol} is not provided in this work.

V836 Tau. A small disk of moderate brightness (with a α_{pol} of 1.8 ± 0.6) is clearly visible in the Q_ϕ image. The apparent disk extent (0.2'') is comparable to ALMA (Long et al. 2019).

MWC480. As discussed in Sects. 4.1.1 and 5.3, this disk is faint given the stellar luminosity but some substructures are immediately visible from the image. Beside the central component and the first ring at 0.6'', a second ring consistent with the disk geometry is marginally visible to the SE at 1'' from the star.

GM Aur. This disk is among the few most spectacular detections of the whole sample. The relatively inclined disk (53° from

Table B.1. Stellar properties of the sample sorted by right ascension. Columns are: target name, Taurus group following [Luhman \(2023\)](#), distance, multiplicity, effective temperature, stellar luminosity, mass, and age.

Target	Group	d (pc)	System	T_{eff} (K)	L_* (L_{\odot})	M_* (M_{\odot})	t (Myr)
CQ Tau	HD35187*	149.4	(*)	6900	6.30 ± 0.27	1.4	12–14
MWC758	HD35187*	155.9	(*)	8200	10.29 ± 0.34	1.7	13–14
RW Aur	L1517	183.3*	(*)(*)	5100+4400	$(1.4 \pm 0.7) \pm 0.3$	1.3 ± 0.9	7.2–9.7
V836 Tau	L1544	167.0	(*)	3600	0.57 ± 0.06	0.4	0.3–0.9
MWC480	L1517	156.2	(*)	8800	19.90 ± 0.65	2.1	6.3–6.6
SU Aur	L1517	157.0	(*)	5800	15.80 ± 2.30	2.5	2.0–2.5
AB Aur	L1517	155.9	(*)	9800	40.58 ± 0.80	2.4	3.9–4.4
GM Aur	L1517	158.1	(*)	4350	1.40 ± 0.18	0.9	1.0–1.6
UY Aur	L1517	152.3	(*)(*)	3800+3600	$(0.65 \pm 0.33) \pm 0.04$	0.5 ± 0.4	0.9–2.1
DS Tau	L1517	158.4	(*)	3900	0.93 ± 0.10	0.5	0.7–1.0
DR Tau	L1558	193.0	(*)	4200	2.53 ± 0.21	0.4	0.3–0.5
DQ Tau	L1558	195.4	(**)	3600	$(0.50 \pm 0.50) \pm 0.04$	0.4	0.7–1.2
EM* LkCa15	L1536	157.2	(*)	4600	1.11 ± 0.11	1.2	3.2–4.5
DO Tau	L1527	138.5	(*)	3800	0.56 ± 0.06	0.5	1.0–1.4
V* V1025 Tau	L1536	167.3	(*)	5800	8.54 ± 1.91	2.0	3.6–5.0
V* HP Tau	L1536	171.2	(*)	5000	1.87 ± 0.61	1.5	3.5–4.1
HQ Tau	L1536	161.4*	(*)	4900	3.24 ± 0.72	1.3	1.5–1.9
DN Tau	L1524 ⁺	128.6	(*)	3850	0.59 ± 0.06	0.5	1.0–1.3
CI Tau	L1536	160.3	(*)	4600	1.35 ± 0.26	1.0	1.3–2.1
DM Tau	L1551	144.0	(*)	3700	0.21 ± 0.03	0.4	2.9–6.0
HN Tau	L1551	134.4*	(*)*	4300	-	(1.6)	-
DL Tau	L1521 ⁺	159.9	(*)	4000	0.89 ± 0.11	0.6	0.9–1.2
GK Tau	L1524	129.1	(*)	4400	0.84 ± 0.06	0.5	0.6–1.0
GI Tau	L1524	129.4	(*)	4100	0.80 ± 0.11	0.7	1.5–1.7
V807 Tau	L1524	184.1*	(**)**	4400+2*3600	$(2.2 \pm 2*0.9) \pm 0.3$	$0.5 \pm 2*0.4$	0.4–0.6
UZ Tau	L1524	123.1*	(**)**	3700+3400	$(0.5 \pm 0.3) \pm 0.09$	0.4 ± 0.3	0.9–1.4
GG Tau	L1551	145.5	(**)(*)	3800+2*3500	$(0.7 \pm 2*0.3) \pm 0.05$	$0.5 \pm 2*0.4$	0.8–2.0
V710 Tau	L1551	146.0	(**)*	3600	0.62 ± 0.05	0.4	0.4–0.8
XZ Tau	L1551	(144)	(*)(*)	2*3600	$2*(0.29 \pm 0.04)$	2*0.4	1.7–2.8
DK Tau	L1524	132.0	(*)(*)	3900+3600	$(1.2 \pm 0.5) \pm 0.05$	0.6 ± 0.4	0.4–0.6
UX Tau	L1551	142.2	(**)**	5200+3700	$(1.8 \pm 0.4) \pm 0.2$	1.4 ± 0.5	6.1–10.0
IQ Tau	L1524	131.5	(*)	3800	0.73 ± 0.13	0.5	0.7–1.0
DH Tau	L1524	133.4	(*)(*)	3600	0.58 ± 0.12	0.4	0.4–0.9
DG Tau	L1524	125.3*	(*)	4000	1.53 ± 0.29	0.6	0.3–0.6
IP Tau	HD28354	129.4	(*)	3800	0.41 ± 0.03	0.6	1.8–2.8
T Tau	T Tau	145.1	(*)(**)	5200	10.40 ± 0.90	$2 \pm 2 \pm 0.5$	0.7–1.0
RY Tau	L1495	138.2*	(*)	5700	11.80 ± 2.90	2.3	2.4–2.6
DE Tau	L1495	128.0	(*)	3700	0.62 ± 0.07	0.4	0.7–1.0
BP Tau	L1495	127.4	(*)	3700	0.69 ± 0.04	0.4	0.5–1.0
V409 Tau	L1524	129.7	(*)	3600	0.22 ± 0.03	0.4	2.7–5.6
CY Tau	L1495	126.3	(*)	3700	0.38 ± 0.04	0.5	1.8–2.2
CX Tau	L1495	126.7	(*)	3500	0.26 ± 0.02	0.4	1.3–2.8
CW Tau	L1495	131.5	(*)	4800	1.59 ± 0.34	1.3	3.1–5.1

Notes. The HD35187 group is formally an association near Taurus. The extended name of the group L1524 is L1524/L1529/B215 ([Luhman 2023](#)) comprising the three individual clouds. The same applies to L1521 (L1521/B213) and L1495 (L1495/B209). In sources with a poor *Gaia* solution (see Sect. 2.1), the distance is accompanied by an asterisk. In all other cases, the tabulated distance has an error of less than 1.5 pc. XZ Tau has no parallax available from *Gaia* and we assume the average distance to its group. Parentheses and asterisks indicate the geometry of disk and star(s), respectively. The effective temperature is relative to the primary and of any companion within 3'' that is bright enough to account for a significant fraction of optical/near-IR photometry (10% of the primary). The masses of T Tau are obtained astrometrically ([Köhler et al. 2016](#)) as most of the observed optical luminosity is from N (Sa and Sb are heavily extincted). The mass of HN Tau is obtained by [Gangi et al. \(2022\)](#) considering obscuration effects.

the ALMA image by [Huang et al. 2020](#)) is visible from both the front and rear disk surfaces that are well separated by a geometrically thick dark lane with an apparent size of nearly $0.5''$. Along the disk major axis, the disk is detected out to $2''$, that is twice as large as the ALMA continuum extent ([Huang et al. 2020](#)). A number of spiral-like structures are visible from the disk front surface as highlighted in Fig. D.1. The ^{12}CO 2–1 line moment-0

map from [Huang et al. \(2021\)](#) shows spiral arms with evident spatial analogies with the SPHERE image.

UY Aur. The polarimetric image shows a number of arc-like structures that are reasonably due to the interaction between the binaries. Given the very low disk extent constrained by ALMA (6 au, [Long et al. 2019](#)), it is probable that none of the detected signal originates from the disk of either component. The Q_{ϕ} im-

Table B.2. Disk properties of the sample sorted by right ascension. Columns are: near-IR, mid-IR, and far-IR excesses relative to the stellar luminosity, approximate disk dust mass, category of the SPHERE image as from Sect. 4.1, measured contrast in scattered light, and reference publication of the SPHERE images shown in this work. **The observing programs are listed in Table C.1.**

Target	NIR (%)	MIR (%)	FIR (%)	M_{dust} (M_{\oplus})	Type	α_{pol}	Reference
CQ Tau	26.8	28.3	24.3	108	Bright	6.4 ± 1.2	Hammond et al. (2022)
MWC758	33.6	9.4	13.7	49	Bright	3.1 ± 0.3	Benisty et al. (2015)
RW Aur	55.1	38.9	18.4	27	Ambient	-	-
V836 Tau	6.4	6.2	4.1	24	Faint	1.8 ± 0.6	-
MWC480	18.7	11.2	5.2	188	Faint	0.5 ± 0.1	-
SU Aur	6.4	5.8	3.7	9	Bright	10.1 ± 0.7	Ginski et al. (2021)
AB Aur	35.6	15.8	17.9	32	Bright	9.7 ± 0.5	Boccaletti et al. (2020)
GM Aur	0	5.7	21.2	130	Bright	16.4 ± 1.8	-
UY Aur	49.6	77.3	53.0	3	Ambient	3.5 ± 0.3	-
DS Tau	14.8	8.2	2.3	12	Faint	< 0.2	-
DR Tau	47.8	44.6	-	128	Bright	3.0 ± 0.6	Mesa et al. (2022)
DQ Tau	14.7	18.3	12.0	79	Faint	0.4 ± 0.2	-
EM* LkCa15	12.1	6.1	7.0	44	Bright	3.7 ± 0.4	Thalmann et al. (2016)
DO Tau	78.0	71.0	56.0	62	Ambient	1.8 ± 0.3	Huang et al. (2022)
V* V1025 Tau	0	0	0.7	1	Faint	< 0.2	-
V* HP Tau	39.3	19.8	-	45	Ambient	-	-
HQ Tau	14.7	10.9	4.2	3	Faint	0.4 ± 0.1	-
DN Tau	9.2	8.9	7.9	41	Faint	0.3 ± 0.3	-
CI Tau	20.7	13.9	9.9	137	Faint	1.4 ± 0.4	Garufi et al. (2022)
DM Tau	0	6.1	14.3	56	Faint	2.6 ± 0.6	-
HN Tau	-	-	-	8	Faint	1.7 ± 1.5	-
DL Tau	10.5	23.0	13.5	130	Faint	0.7 ± 0.4	-
GK Tau	30.6	22.0	8.4	2	Faint	0.3 ± 0.3	-
GI Tau	13.3	17.5	-	6	Faint	0.4 ± 0.3	-
V807 Tau	4.8	4.2	1.6	17	Faint	< 0.3	-
UZ Tau	26.0	-	-	14	Faint	2.5 ± 0.4	Zurlo et al. in prep.
GG Tau	17.6	14.8	12.9	295	Bright	8.7 ± 1.2	Keppeler et al. (2020)
V710 Tau	15.9	10.1	3.6	34	Faint	2.7 ± 0.4	-
XZ Tau	102	120	55.0	2	Ambient	-	Zurlo et al. in prep.
DK Tau	16.1	12.2	6.4	9	Faint	< 0.2	-
UX Tau	8.8	3.7	8.4	33	Ambient	8.7 ± 0.9	Ménard et al. (2020)
IQ Tau	20.5	10.8	6.1	32	Faint	1.7 ± 0.7	-
DH Tau	17.6	6.1	5.6	1	Faint	0.9 ± 0.2	van Holstein et al. (2021)
DG Tau	23.2	53.0	57.1	162	Ambient	6.5 ± 0.8	-
IP Tau	14.0	8.8	5.3	4	Faint	1.3 ± 0.2	-
T Tau	27.9	55.4	56.0	67	Ambient	-	-
RY Tau	22.9	21.9	12.0	52	Ambient	3.8 ± 0.6	Valegård et al. (2022)
DE Tau	15.7	11.0	8.3	15	Faint	0.6 ± 0.3	-
BP Tau	13.7	9.8	5.6	20	Faint	0.5 ± 0.3	-
V409 Tau	7.5	15.6	-	9	Faint	2.9 ± 0.5	-
CY Tau	11.3	6.0	3.1	43	Faint	0.6 ± 0.1	-
CX Tau	10.6	13.3	7.1	5	Faint	0.4 ± 0.3	-
CW Tau	25.0	19.0	6.9	27	Faint	1.5 ± 0.3	-

Notes. Unreported excesses are due to uncertainty in the SED or poor available photometry. Unreported contrasts are due to the presence of ambient material that biases the measurement. All SPHERE datasets are in the H band except CQ Tau (J band), MWC758 (K), LkCa15 (J), and CY Tau (K).

age shows some strong negative signal due to the complex summation of the individual stellar contributions.

DS Tau. No *QU* pattern is visible. Nonetheless, some Q_{ϕ} signal that is spatially consistent with the ALMA image is visible from Fig. 6. A SPHERE image at longer wavelengths (K band) reveals a relatively clear detection (Ren et al. in prep.). Yet, in this work this is treated as a non-detection.

DQ Tau. A relatively clear disk is detected out to $0.2''$, that is the extent of the ALMA continuum emission by Long et al. (2019). Nonetheless, this disk is rather massive in dust ($79 M_{\oplus}$) making it a good example of a self-shadowed disk (see Fig. 4).

V V1025 Tau.* No *QU* pattern is visible. It is treated as a non-detection. As discussed throughout the text and shown in Fig. 4, it is a Class III object. It is very close in space to HP Tau (see next source).

V HP Tau.* A very complex pattern of arc-like structures is visible from the Q_{ϕ} image. The brightest arcs are distributed within the inner $2''$ but some fainter structures are visible out to the detector edge at $6''$. The disk has an appreciable dust mass ($45 M_{\oplus}$ from our calculation). No companion that may be responsible for the peculiar features observed is visible from the SPHERE image, but the source is close to CoKu HP Tau G3

Table C.1. Details of observations.

Target	Program (PI)	Filter	Observing night	DIT (sec)	NDIT	t_{int} (min)
CQ Tau	098.C-0760(B) (Benisty)	J	2017-10-06	50	40	33
MWC758	106.21HJ.001 (Benisty)	K _s	2020-12-25	16	140	37
RW Aur	0104.C-0122(A) (Facchini)	H	2019-12-04	16	192	51
V836 Tau	0102.C-0453(A) (GTO, Beuzit)	H	2019-09-01	32	20	11
MWC480	0101.C-0867(A) (Keppler)	H	2018-11-17	32	108	57
SU Aur	1104.C-0415(E) (DESTINYs, Ginski)	H	2019-12-14	32	104	55
AB Aur	0104.C-0157(B) (Boccaletti)	H	2019-12-18	32	176	94
GM Aur	0100.C-0452(A) (Benisty)	H	2018-01-02	64	32	34
UY Aur	1104.C-0415(G) (DESTINYs, Ginski)	H	2021-01-21	16	94	25
DS Tau	0102.C-0453(A) (GTO, Beuzit)	H	2018-12-25	32	20	11
DR Tau	0102.C-0453(A) (GTO, Beuzit)	H	2018-11-25	32	20	11
DQ Tau	1104.C-0415(E) (DESTINYs, Ginski)	H	2019-11-24	64	56	60
EM* LkCa15	096.C-0248(A) (GTO, Beuzit)	J	2020-12-07	32	120	64
DO Tau	1104.C-0415(E) (DESTINYs, Ginski)	H	2019-12-19	64	56	60
V* V1025 Tau	1104.C-0415(A) (DESTINYs, Ginski)	H	2019-12-15	64	56	60
V* HP Tau	1104.C-0415(E) (DESTINYs, Ginski)	H	2019-12-19	64	56	60
HQ Tau	0102.C-0453(A) (GTO, Beuzit)	H	2018-11-28	32	20	11
DN Tau	0102.C-0453(A) (GTO, Beuzit)	H	2019-08-15	32	20	11
CI Tau	0100.C-0452(A) (Benisty)	H	2018-01-01	64	32	34
DM Tau	0101.C-0867(A) (Keppler)	H	2018-10-01	96	36	58
HN Tau	1104.C-0415(E) (DESTINYs, Ginski)	H	2019-12-13	64	12	13
DL Tau	0102.C-0453(A) (GTO, Beuzit)	H	2018-12-28	32	20	11
GK Tau	0102.C-0453(A) (GTO, Beuzit)	H	2019-10-24	32	20	11
GI Tau	0102.C-0453(A) (GTO, Beuzit)	H	2018-11-26	32	20	11
V807 Tau	1104.C-0415(E) (DESTINYs, Ginski)	H	2019-12-30	0.837	4320	60
UZ Tau	0100.C-0408(B) (Cieza)	H	2017-10-27	64	108	115
GG Tau	198.C-0209(N) (Beuzit)	H	2016-11-18	4	660	44
V710 Tau	1104.C-0415(A) (DESTINYs, Ginski)	H	2019-12-20	4	1536	102
XZ Tau	0100.C-0408(A) (Cieza)	H	2017-10-12	64	90	96
DK Tau	0102.C-0453(A) (GTO, Beuzit)	H	2019-09-01	32	20	23
UX Tau	0100.C-0452(B) (Benisty)	H	2017-10-05	64	48	51
IQ Tau	0102.C-0453(A) (GTO, Beuzit)	H	2018-12-18	32	44	23
DH Tau	0102.C-0916(A) (GTO, Beuzit)	H	2019-10-23	64	36	38
DG Tau	1104.C-0415(D) (DESTINYs, Ginski)	H	2021-10-28	64	56	60
IP Tau	1104.C-0415(A) (DESTINYs, Ginski)	H	2019-12-14	64	56	60
T Tau	198.C-0209(N) (GTO, Beuzit)	H	2016-11-18	16	64	17
RY Tau	1104.C-0415(E) (DESTINYs, Ginski)	H	2018-12-16	32	68	36
DE Tau	0102.C-0453(A) (GTO, Beuzit)	H	2019-10-11	32	20	11
BP Tau	0102.C-0453(A) (GTO, Beuzit)	H	2018-12-22	32	20	11
V409 Tau	1104.C-0415(A) (DESTINYs, Ginski)	H	2019-11-16	64	56	60
CY Tau	108.22EE.001 (Benisty)	K _s	2021-12-26	32	96	51
CX Tau	0102.C-0453(A) (GTO, Beuzit)	H	2019-10-23	32	20	11
CW Tau	0102.C-0453(A) (GTO, Beuzit)	H	2018-12-22	32	32	17

Notes. DIT is the detector integration time, NDIT is the number of detector integration times, t_{int} is the total on-target integration time ($t_{\text{obs}} = \text{DIT} \times \text{NDIT}$).

(17'') and V1025 Tau (21'', see previous source), as well as to the bright reflection nebula GN 04.32.8.

HQ Tau. A very faint ($\alpha_{\text{pol}}=0.4\pm0.1$) and compact (0.2'') *QU* pattern is visible, suggesting a vertically oriented disk that is consistent with the ALMA image by Long et al. (2019). With only 3 M_{\oplus} estimated, this is one of the lowest mass disks ever resolved in the near-IR (see Fig. 4).

DN Tau. Despite the very low contrast measured (0.3 ± 0.3), a faint *QU* pattern is visible out to 0.6''. As shown in Fig. 6, it is a prototypical example of self-shadowed disk.

DM Tau. The detection of the faint disk of DM Tau is described in depth in Sect. 5.2.

HN Tau. The AO correction is imperfect because of the low stellar luminosity ($G=13.4$, see Appendix B). It is treated as a

marginal detection because the Q_{ϕ} signal aligns with the disk position angle by Long et al. (2019). However, the calculated α_{pol} and relative error (1.7 ± 1.5) reflects this uncertainty.

DL Tau. Despite the low contrast measured (0.7 ± 0.4), the disk is resolved out to 0.8''. As shown in Fig. 6, it is a prototypical example of self-shadowed disk where some disk substructures are visible after careful inspection.

GK Tau. No *QU* pattern is visible. It is treated as a non-detection. The low dust mass calculated (2 M_{\oplus}) suggests a small disk (<9 au in size from Long et al. 2019).

GI Tau. A very faint ($\alpha_{\text{pol}}=0.4\pm0.3$) and compact (0.2'') *QU* pattern is visible. Similarly to HQ Tau, with only 6 M_{\oplus} estimated, this is one of the lowest mass disks ever resolved in the near-IR.

Table D.1. Disk geometry interpreted from the SPHERE image.

Target	r_{out} (")	i	P.A. (°)
CQ Tau	0.6	Intermediate	45
MWC758	0.6	Mild	-
RW Aur	0.5	Intermediate	45
V836 Tau	0.2	Intermediate	120
MWC480	1.0	Intermediate	50
SU Aur	0.5	Mild	-
AB Aur	2.4	Intermediate	45
GM Aur	2.0	Intermediate	60
UY Aur	-	-	-
DS Tau	-	-	-
DR Tau	0.7	-	-
DQ Tau	0.2	Intermediate	20
LkCa15	0.7	Intermediate	60
DO Tau	0.6	-	-
V1025 Tau	-	-	-
HP Tau	-	-	-
HQ Tau	0.2	-	-
DN Tau	0.6	-	-
CI Tau	0.5	Intermediate	10
DM Tau	0.4	Intermediate	90
HN Tau	-	-	-
DL Tau	0.8	Intermediate	50
GK Tau	-	-	-
GI Tau	0.2	-	-
V807 Tau	-	-	-
UZ Tau	0.6	High	90
GG Tau	1.8	Intermediate	110
V710 Tau	0.4	Intermediate	70
XZ Tau	-	-	-
DK Tau	-	-	-
UX Tau	0.6	Intermediate	80
IQ Tau	0.5	High	45
DH Tau	0.2	-	-
DG Tau	0.7	Intermediate	140
IP Tau	0.3	Intermediate	70
T Tau	-	-	-
RY Tau	-	-	-
DE Tau	0.2	-	-
BP Tau	0.3	Intermediate	150
V409 Tau	0.5	High	40
CY Tau	0.2	-	-
CX Tau	0.4	-	-
CW Tau	0.4	High	70

Notes. Columns are: target, outermost separation with detectable signal, indicative inclination and position angle.

V807 Tau. The SPHERE image is non-coronagraphic. The binary companion Bab is visible from the image at 0.17" and is in turn marginally resolved. Owing to the presence of the companion, no *QU* pattern can be identified. It is treated as a non-detection. The ALMA image by [Akeson et al. \(2019\)](#) reveals a marginally resolved emission where the individual components of A and Bab are not resolved.

UZ Tau. A clear, moderately inclined disk is visible from the Q_ϕ image. The measured contrast (2.5 ± 0.4) is close to the formal limit between bright and faint disks adopted in this work. As shown in Fig. 6, it is a prototypical example of self-shadowed disk. This source will be studied in a forthcoming publication ([Zurlo et al.](#))

V710 Tau. The clear disk detected in the Q_ϕ image has a dip with negative values to the North. Along the disk major axis, the disk appears much more extended to the East (0.4") than to the West (0.25"). We are prone to consider these effects as real, and not due to an imperfect centering or stellar polarization subtraction.

XZ Tau. Multiple, extended structures that are at least partly due to the interaction between the binaries are visible in the polarimetric image. This source will be studied in a forthcoming publication ([Zurlo et al.](#))

DK Tau. No *QU* pattern is visible. It is treated as a non-detection. The low dust mass calculated ($9 M_\oplus$) suggests a small disk (<15 au in size from [Long et al. 2019](#)).

IQ Tau. A faint but yet extended (0.5") disk is visible from the Q_ϕ image, with an extent comparable to ALMA (see Fig. 6).

DG Tau. The Q_ϕ image of DG Tau reveals a complicated field, with a prominent contribution from the natal envelope and from extended arms that will be studied in a forthcoming publication.

IP Tau. The detection of the faint disk of IP Tau is described in depth in Sect. 5.2.

T Tau. In line with previous work (e.g., [Kasper et al. 2020](#)), the Q_ϕ image of T Tau reveals an extremely complicated field, with multiple arms, arcs, and dips that will be studied in a forthcoming publication ([van Holstein et al.](#)).

DE Tau. The Q_ϕ image only reveals faint signal along the SE-NW axis from the inner 0.15".

BP Tau. Similarly to V710 Tau, the disk that is clearly visible from the Q_ϕ image exhibits a dip to the North and an obvious azimuthal asymmetry, with the SE portion of the disk being brighter than the NW region.

V409 Tau. A clear, inclined disk is visible from the Q_ϕ image in line with the ALMA image ($i = 69^\circ$, [Long et al. 2019](#)). The measured contrast (2.9 ± 0.5) is at the formal limit between bright and faint disks adopted in this work. However, it is known that inclined disks appear artificially brighter in scattered light because we inevitably probe the forward peak of the phase function ([Garufi et al. 2022](#)).

CY Tau. The Q_ϕ image reveals a faint emission with an approximate position angle of 160° pointing to a relatively inclined disk nearly vertically oriented. This possible morphology is coarsely consistent with the archival ALMA data from program 2013.1.00498 (PI: Perez, L.).

CX Tau. The presence of a faint *QU* pattern can be inferred out to 0.35". The presence of a gaseous disk more extended than 75 au ([Facchini et al. 2019](#)) surrounding a compact continuum emission (14 au) indicates that this is also a self-shadowed disk.

CW Tau. A clearly inclined disk is inferred from the Q_ϕ image that is geometrically consistent with the ALMA image by [Ueda et al. \(2022\)](#).




Fig. B.1. Spectral energy distributions of the entire sample. The de-reddened literature photometry (see Sect. 2.3) is indicated by the red symbols while the *Phoenix* model for the stellar photosphere by the black line.




Fig. D.1. Multi-component view of GM Aur. *Top-left:* SPHERE image from this work obtained with an adaptive de-noising procedure based on the adaptive kernel smoothing technique and implemented in the SPLASH software (Price 2007). *Top-right:* ALMA continuum image in Band 4 from Huang et al. (2020). *Bottom-left:* ALMA ^{12}CO 2–1 line moment-0 map from Huang et al. (2021). *Bottom-right:* Combined view. The most obvious arms are highlighted in this panel. The common spatial scale is indicated to the bottom right.