Summary Information Sheet

Comets

A comet's nucleus is a "dirty snowball" consisting of frozen gases (ices) and dust. Orbital periods range from about 3 years to millions of years and some comets move in highly inclined orbits. Close to the Sun, ices evaporate and dust grains are released from the nucleus, forming a "coma" (the comet's head) and one or more tails. Some comets become very bright and have spectacular tails.




Sunlit side




The Bayeux Tapestry with Comet Halley (1066)

Dust grains: organic compounds and silicates

Frozen gases (ices): mainly water, carbon monoxide, carbon dioxide



Orbit and evolution of a comet




Internal structure is loose and fluffy some comets break up Comets carry material dating from the formation of the Solar System


Dust tail up to 100 mio km long or more



Very dark surface layer reflects only 3-4% of incoming sunlight



Comet tail evolution (Halley 1910)



Tails of comet Hale-Bopp (1998)



Comet Halley's nucleus (ESA Giotto, 1986)



Comet Borrelly (NASA Deep Space 1, 2001)



Broken-up comet SL-9 hits Jupiter (1994)

Physical Data

Property Distance from the Sun Orbital period **Dimensions** Mass Density

Comet Halley

88-5300 mio km 76 years Nucleus: 16 x 8 x 7 km $8 \times 10^{13} \text{ kg}$ $100 \, \text{kg/m}^3$

| Asteroid Gaspra |
|-----------------|
| 332 mio km |
| 3.29 years |
| 19 x 12 x 11 km |
| ? |
| ? |

Mars moon Deimos

1.26 days 15 x 12 x 11 km $1.8 \times 10^{15} \text{ kg}$ 1700 kg/m^3

Concept: Bernhard Mackowiak

Images from ESO, ESA or NASA except otherwise stated

For comparison